Information For the Use of
Metoprolol Succinate (100mg) PR Tablets

Metoperil XL 100 Tablet

Metoperil XL 100 Tablet

Composition:

Each prolonged-release tablet contains:

- **Metoprolol Succinate** – 100 mg

Pharmacological Action:

Metoprolol succinate is a **selective β1-adrenergic receptor blocker** that reduces the effects of sympathetic stimulation on the heart.

- **Heart Rate Reduction:** Slows the heart rate by reducing β1-adrenergic stimulation.
- **Reduced Cardiac Workload:** Decreases myocardial contractility and cardiac oxygen demand.
- **Blood Pressure Control:** Reduces cardiac output and suppresses sympathetic activity, helping lower blood pressure.
- **Antianginal Action:** Reduces myocardial oxygen demand and helps prevent anginal symptoms.
- **Cardioprotective Effect:** In appropriately selected patients with heart failure, reduces harmful sympathetic activation and improves clinical outcomes.

Indications:

Metoperil XL 100 Tablet is indicated for the management of:

- **Hypertension**
- **Angina Pectoris**
- **Chronic Heart Failure with Reduced Ejection Fraction (HFrEF)** in clinically stable patients
- **Rate Control in Certain Tachyarrhythmias**
- Prevention of recurrent myocardial infarction, where clinically appropriate

---

Dosage and Administration:

- **Recommended Dose:** Dose is individualized according to the indication and patient response. **100 mg once daily** may be used in appropriate patients, or as directed by the physician.
- Take once daily, preferably at the same time each day.
- Swallow the prolonged-release tablet **whole with water**.
- Do not crush or chew the tablet, as this may interfere with the prolonged-release characteristics.
- It may be taken with or immediately after food.

- Dose titration should be performed gradually under medical supervision.
- **Do not discontinue abruptly**, particularly in patients with coronary artery disease, because sudden withdrawal may worsen angina or precipitate serious cardiac events.

Precautions:

- Monitor **heart rate and blood pressure** during treatment.
- Use cautiously in patients with **bradycardia, conduction disorders, peripheral vascular disease, diabetes, or thyroid disorders**.
- Metoprolol may mask some symptoms of **hypoglycemia**, particularly tachycardia.
- Use cautiously in patients with asthma or COPD; although β1-selective, bronchospasm may occur, particularly at higher doses.
- Patients with heart failure should be clinically stable before initiation and dose escalation.
- If severe dizziness, fainting, unusually slow heartbeat, or worsening breathlessness occurs, seek medical advice.
- **Pregnancy and Lactation:** Use only when clearly indicated and under medical supervision.

Contraindications:

- Known hypersensitivity to **metoprolol or any component of the formulation**.
- **Severe bradycardia**.
- Second- or third-degree **heart block** in patients without a functioning pacemaker.
- **Sick sinus syndrome** without a functioning pacemaker.
- Cardiogenic shock.
- **Decompensated heart failure** requiring intravenous inotropic support.
- Clinically significant hypotension.

Adverse Reactions:

Common:

- Fatigue
- Dizziness
- Headache
- Slow heart rate
- Cold hands and feet
- Gastrointestinal discomfort
- Sleep disturbances

Less Common but Serious:

- Excessive bradycardia

- Hypotension
- Heart block
- Worsening heart failure
- Bronchospasm
- Peripheral circulatory disturbances
- Severe hypersensitivity reactions

Seek medical attention for fainting, severe shortness of breath, chest pain, or markedly slow heartbeat.

Drug Interactions:

- **Verapamil and diltiazem** may increase the risk of bradycardia, hypotension, and heart block.
- Other **beta-blockers** may produce additive cardiovascular effects.
- **Digoxin** may increase the risk of bradycardia and conduction abnormalities.
- **Clonidine:** Concomitant use requires careful management; abrupt withdrawal of clonidine while taking a beta-blocker can cause significant rebound hypertension.
- **Insulin and oral antidiabetic medicines:** Metoprolol may mask some symptoms of hypoglycemia.
- Other medicines that lower blood pressure or heart rate may produce additive effects.
- Certain medicines affecting **CYP2D6**, such as some antidepressants, may increase metoprolol exposure.


Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture, direct sunlight, and excessive heat.
- Keep tablets in the original packaging.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. **Metoperil XL 100 Tablet should be used under the supervision of a registered medical practitioner**, with dosage individualized according to the indication, blood pressure, heart rate, and clinical response.

Important: Metoprolol succinate prolonged-release tablets should be **swallowed whole and should not be crushed or chewed**. Do not stop treatment abruptly without medical advice, particularly in patients with coronary artery disease.



Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for