Metrohype 600 Tablet

METROHYPE 600 TABLET

Metronidazole 600 mg Tablet

1. Composition

Each uncoated tablet contains:

Metronidazole IP — 600 mg
Excipients — q.s.

Verify the exact approved formulation and pharmacopoeial specification before printing.

2. Pharmacological Action

Metronidazole is a nitroimidazole antimicrobial agent. It acts against susceptible anaerobic bacteria and certain protozoa by interfering with microbial DNA, thereby inhibiting growth and causing cell death.

3. Indications

Metronidazole may be prescribed for susceptible infections, including:

- Amoebiasis
- Giardiasis
- Trichomoniasis
- Anaerobic bacterial infections
- Bacterial vaginosis, where appropriate

Use should be based on the confirmed or suspected organism and applicable treatment guidelines.

4. Dosage and Administration

Administer orally as directed by the treating physician. The dose, frequency and duration depend on the type and severity of infection, patient age, clinical condition and applicable treatment guidelines.

Swallow with water; taking the tablet with or after food may help reduce stomach discomfort. Complete the prescribed course.

5. Precautions

- Use cautiously in patients with liver disease or a history of neurological disorders.
- Prolonged or repeated treatment may require clinical and laboratory monitoring.
- Avoid alcohol during treatment and for at least 48 hours after the last dose, as a reaction may occur.
- Seek medical advice if numbness, tingling, unsteadiness or unusual neurological symptoms develop.

6. Contraindications

Do not use in patients with known hypersensitivity to metronidazole or other nitroimidazole derivatives. Check the current approved product information for additional contraindications.

7. Adverse Reactions

Possible adverse effects include:

- Nausea, vomiting or abdominal discomfort
- Diarrhoea or loss of appetite
- Metallic taste or dry mouth
- Headache or dizziness
- Darkened urine

Rare but serious reactions may include allergic reactions, seizures, or peripheral neuropathy. Seek urgent medical attention for severe symptoms.

8. Drug Interactions

Metronidazole may interact with:

- **Warfarin and other coumarin anticoagulants:** may increase anticoagulant effects.
- **Lithium:** may increase lithium concentrations.
- **Disulfiram:** concurrent use may cause adverse neuropsychiatric effects.
- **Certain enzyme-inducing or enzyme-inhibiting medicines:** may alter metronidazole exposure.

Inform the prescribing physician about all medicines and supplements being used.

9. Pregnancy and Lactation

Use during pregnancy or breastfeeding only after an appropriate clinical assessment and in accordance with current prescribing guidance.

10. Storage

Store below 30°C, protected from moisture and direct sunlight, unless the approved product label specifies otherwise. Keep out of reach of children.

11. Note

For oral use only. To be sold by retail on the prescription of a Registered Medical Practitioner, where required by applicable regulations.

Important: This is a draft product-literature template, not an official approved package insert. Confirm all claims, dosage guidance, warnings and legal wording against the current approved prescribing information for Metrohype 600 Tablet before printing or distribution.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for