Information For the Use of
Minocycline (50mg) Tablets

Minoclub 50 Tablet

MINOCLUB 50 TABLET

Minocycline Hydrochloride (50 mg)

1. Composition

Each film-coated tablet contains:
Minocycline Hydrochloride equivalent to Minocycline 50 mg
Excipients q.s.

2. Pharmacological Action

Minocycline is a **tetracycline-class antibiotic**. It inhibits bacterial protein synthesis by binding to the 30S ribosomal subunit, thereby preventing the attachment of aminoacyl-tRNA to the bacterial ribosome and inhibiting bacterial growth.

3. Indications

Minocycline is indicated for the treatment of susceptible bacterial infections, including:

- **Acne vulgaris** and inflammatory acne.
- Certain respiratory tract infections caused by susceptible organisms.
- Selected skin and soft-tissue infections.
- Certain genitourinary infections caused by susceptible bacteria.
- Other infections where the causative organism is susceptible to minocycline.

Antibiotic selection should be guided by the suspected/confirmed pathogen and local susceptibility patterns.

4. Dosage and Administration

Adults: The dosage depends on the indication and severity of infection. For acne, commonly used regimens may include **50 mg once or twice daily**, as directed by the physician.

The tablet should be swallowed whole with a sufficient quantity of water. It may be taken with or without food, although taking it with food may reduce gastrointestinal irritation.

Avoid taking minocycline simultaneously with **antacids containing aluminium, calcium or magnesium, iron preparations, or other mineral supplements**, as these may reduce absorption.

5. Precautions

- Use cautiously in patients with **hepatic impairment**.
- Minocycline may cause dizziness, light-headedness or vertigo; patients should exercise caution while driving or operating machinery.
- Prolonged therapy may require appropriate clinical monitoring.
- Tetracyclines may cause **photosensitivity**; excessive exposure to sunlight should be avoided.
- Minocycline may cause pigmentation of the skin, mucous membranes, nails or teeth, particularly with prolonged use.
- Use during tooth development may cause permanent tooth discoloration.
- Discontinue and seek medical attention if severe headache, visual disturbances, significant skin reactions or symptoms suggestive of liver injury occur.
- As with other antibiotics, prolonged use may result in overgrowth of non-susceptible organisms.

6. Contraindications

Minocycline is contraindicated in:

- Patients with known hypersensitivity to **minocycline, tetracyclines**, or any component of the formulation.
- **Pregnancy**, particularly because of potential effects on fetal bone and teeth.
- **Breastfeeding**, unless specifically advised by the physician.
- Children below **8 years of age**, except where specifically indicated by a physician.

7. Adverse Reactions

Possible adverse reactions include:

- Nausea, vomiting and diarrhoea
- Abdominal discomfort
- Dizziness or vertigo
- Headache
- Skin rash and photosensitivity
- Changes in skin, teeth or mucosal pigmentation

Rare but serious reactions may include **hepatotoxicity, intracranial hypertension, severe hypersensitivity reactions, autoimmune reactions and serious skin reactions**.

8. Drug Interactions

Minocycline absorption may be reduced by:

- **Aluminium-, calcium- or magnesium-containing antacids**
- Iron-containing preparations
- Calcium supplements and other mineral-containing products

Other important interactions may occur with anticoagulants, isotretinoin and certain medicines affecting hepatic function. The physician should be informed about all prescription, OTC and herbal medicines being used.

9. Storage

Store below **30°C**, protected from light and moisture.

Keep out of reach of children.

10. Note / Important

Minocycline is an antibiotic intended for **susceptible bacterial infections** and should not be used for viral infections.

Take the medicine exactly as prescribed and complete the recommended course unless otherwise directed by the physician.

For use under medical supervision.

Keep out of reach of children.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for