Milritype 10 Injection

MILRITYPE 10 INJECTION

Milrinone Injection 10 mg

COMPOSITION

Each ampoule/vial contains:

Milrinone Lactate equivalent to Milrinone 10 mg
Suitable Sterile Vehicle q.s.

Confirm the salt form, total ampoule volume and strength per mL against the approved product specification before printing.

PHARMACOLOGICAL ACTION

Milrinone is a phosphodiesterase-3 (PDE-3) inhibitor with positive inotropic and vasodilator properties. It increases intracellular cyclic adenosine monophosphate (cAMP), enhancing myocardial contractility while relaxing vascular smooth muscle.

It increases cardiac output and reduces systemic and pulmonary vascular resistance. Its combined effects help improve cardiac performance in selected patients with acute decompensated heart failure.

INDICATIONS

Milritype 10 Injection may be indicated for:

- Short-term intravenous treatment of acute decompensated heart failure.
- Low cardiac output associated with impaired myocardial contractility.
- Selected cases requiring temporary inotropic support in intensive care.
- Specialist-directed haemodynamic support following cardiac surgery, where clinically appropriate.

DOSAGE AND ADMINISTRATION

Milrinone must be administered by **controlled intravenous infusion under continuous medical supervision**.

Commonly referenced adult dosing:

- **Loading dose:** 50 mcg/kg administered slowly over 10 minutes, when appropriate.
- **Maintenance infusion:** Usually 0.375–0.75 mcg/kg/min, adjusted according to haemodynamic response and tolerability.
- The loading dose may be omitted in patients at increased risk of hypotension or when clinically inappropriate.
- Dosage reduction is required in renal impairment because milrinone is primarily eliminated by the kidneys.

Administration precautions:

- Use an infusion pump and the dilution specified in the approved product information.
- Monitor blood pressure, ECG, heart rate, fluid balance, renal function and electrolytes.
- Avoid rapid administration of the loading dose.
- The dose and duration must be individualized by the treating specialist.

PRECAUTIONS

- Milrinone may cause hypotension due to peripheral vasodilation.
- Monitor continuously for ventricular and supraventricular arrhythmias.
- Use cautiously in patients with low blood pressure, myocardial ischaemia or a history of significant arrhythmias.
- Assess and correct hypokalaemia and other electrolyte disturbances.
- Reduce the infusion rate in renal impairment.
- Monitor platelet counts when clinically indicated.
- Prolonged use beyond the recommended short-term period requires careful specialist assessment.
- Milrinone does not replace the treatment of the underlying cause of heart failure.

CONTRAINDICATIONS

- Hypersensitivity to milrinone or any formulation component.
- Severe uncorrected hypovolaemia.
- Severe obstructive aortic or pulmonary valvular disease, where increased cardiac output may be inappropriate.
- Other contraindications specified in the approved product information.

ADVERSE REACTIONS

Common

- Hypotension
- Headache
- Nausea
- Injection-site reactions

- Changes in heart rate

Serious / Less Common

- Ventricular arrhythmias
- Supraventricular arrhythmias
- Severe hypotension
- Myocardial ischaemia
- Thrombocytopenia
- Rare hypersensitivity reactions

DRUG INTERACTIONS

- **Other vasodilators:** May increase the risk of hypotension.
- **Diuretics:** Electrolyte disturbances may increase arrhythmia risk.
- **Other inotropic agents:** May produce additive cardiovascular effects.
- **Antihypertensive medicines:** May increase the risk of low blood pressure.
- Medicines that affect cardiac rhythm should be reviewed before administration.

USE IN PREGNANCY AND LACTATION

Use during pregnancy only when clearly indicated and when the expected benefit justifies the potential risk.

Information regarding use during lactation is limited. Treatment should be guided by a qualified specialist.

STORAGE

- Store according to the approved product label.
- Protect from excessive heat and light.
- Do not freeze unless specifically permitted by the manufacturer.
- Inspect the solution before use; do not use if it contains visible particles or unexpected discoloration.
- Keep out of reach of children.

NOTE

Milritype 10 Injection is intended for **hospital use under qualified medical supervision**. The exact strength, dilution, infusion rate and approved indications must be verified against the registered Indian product information before final printing.

For professional use only.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for