Information For the Use of
Naproxen (500mg) Tablets

Naprohike 500 Tablet

Naprohike 500 Tablet

Composition:

Each tablet contains:

- **Naproxen** – 500 mg

Pharmacological Action:

Naproxen is a **non-steroidal anti-inflammatory drug (NSAID)** with analgesic, anti-inflammatory, and antipyretic properties.

- **Anti-inflammatory Action:** Inhibits cyclooxygenase (COX) enzymes, reducing prostaglandin synthesis and thereby decreasing inflammation.
- **Analgesic Action:** Reduces prostaglandin-mediated sensitization of pain receptors, helping relieve mild to moderate pain.
- **Antipyretic Action:** Reduction of prostaglandin synthesis in the central nervous system helps lower fever.
- **Prolonged Effect:** Provides relatively long-lasting relief due to its longer duration of action compared with some other NSAIDs.

Indications:

Naprohike 500 Tablet is indicated for the management of:

- **Rheumatoid Arthritis**
- **Osteoarthritis**
- **Ankylosing Spondylitis**
- **Acute Musculoskeletal Pain**
- **Acute Gout**
- **Dysmenorrhea**
- **Tendonitis and Bursitis**
- Other painful inflammatory conditions as prescribed by the physician

---

Dosage and Administration:

- **Recommended Dose:** Dose varies according to the indication. **500 mg once or twice daily** may be used in appropriate adults, as directed by the physician.

- Take with food or milk if gastrointestinal discomfort occurs.
- Swallow the tablet whole with water.
- Use the **lowest effective dose for the shortest duration** necessary.
- Avoid taking more than the prescribed dose.
- Do not combine with other NSAIDs unless specifically advised by the physician.

Precautions:

- Use cautiously in patients with a history of **peptic ulcer disease, gastrointestinal bleeding, hypertension, heart failure, kidney disease, or liver disease**.
- NSAIDs may increase the risk of **gastrointestinal bleeding, cardiovascular events, and renal impairment**, particularly with prolonged use or in high-risk patients.
- Elderly patients may have an increased risk of gastrointestinal, renal, and cardiovascular adverse effects.
- Maintain adequate hydration, particularly in patients at risk of renal impairment.
- Avoid prolonged self-medication.
- **Pregnancy:** Avoid use during the **third trimester**; use during pregnancy only when specifically advised by a physician.
- **Lactation:** Use only when considered necessary by the physician.

Contraindications:

- Known hypersensitivity to **naproxen, other NSAIDs, or any component of the formulation**.
- History of **aspirin/NSAID-induced asthma, urticaria, or allergic reactions**.
- Active or recurrent **peptic ulcer or gastrointestinal bleeding**.
- Severe renal or hepatic impairment.
- Severe heart failure.
- Use for **peri-operative pain in the setting of coronary artery bypass graft (CABG) surgery**.
- Third trimester of pregnancy.

Adverse Reactions:

Common:

- Nausea
- Dyspepsia
- Heartburn
- Abdominal pain
- Constipation or diarrhea
- Headache
- Dizziness

Less Common but Serious:

- Gastrointestinal ulceration or bleeding
- Kidney impairment
- Liver injury
- Fluid retention and edema
- Increased blood pressure
- Serious cardiovascular events
- Severe skin reactions
- Anaphylactic or other serious hypersensitivity reactions

Seek immediate medical attention for **black stools, vomiting blood, severe abdominal pain, chest pain, difficulty breathing, facial swelling, or markedly reduced urine output**.

Drug Interactions:

- **Anticoagulants** such as warfarin may significantly increase the risk of gastrointestinal bleeding.
- **Antiplatelet medicines**, including aspirin and clopidogrel, may increase bleeding risk.
- **Other NSAIDs** may increase gastrointestinal and renal toxicity and should generally be avoided concurrently.
- **ACE inhibitors/ARBs and diuretics** may increase the risk of renal impairment and may reduce antihypertensive efficacy.
- **Lithium:** Naproxen may increase lithium concentrations and toxicity risk.
- **Methotrexate:** NSAIDs may increase methotrexate exposure and toxicity.
- **Cyclosporine or tacrolimus:** May increase the risk of nephrotoxicity.
- **Corticosteroids and SSRIs/SNRIs** may increase the risk of gastrointestinal bleeding.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture, direct sunlight, and excessive heat.
- Keep tablets in the original packaging.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. **Naprohike 500 Tablet should be used under the supervision of a registered medical practitioner.** Use the lowest effective dose for the shortest appropriate duration to minimize NSAID-related gastrointestinal, renal, and cardiovascular risks.

Important: Avoid concurrent use with other NSAIDs unless specifically prescribed. Patients taking **warfarin or other anticoagulants/antiplatelet medicines** should consult their physician before using naproxen because of the increased risk of bleeding.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for