Information For the Use of
Nebivolol 2.5 mg Tablet

Nebiroll 2.5 Tablet

Brand Name: Nebiroll 2.5 Tablet
Dosage Form: Tablet
Composition:
Each tablet contains:
• Nebivolol Hydrochloride IP equivalent to Nebivolol 2.5 mg

Description:
Nebiroll 2.5 Tablet contains Nebivolol, a highly selective β₁-adrenergic receptor blocker with nitric oxide-mediated vasodilatory properties. It is indicated for the management of essential hypertension and may also be used in selected patients with stable chronic heart failure as advised by the physician. Nebivolol helps reduce blood pressure by lowering heart rate, decreasing myocardial oxygen demand, and promoting vasodilation, thereby improving cardiovascular function.

Indications:
Indicated for the treatment of:
• Essential hypertension
• Stable chronic heart failure (as an adjunct to standard therapy in appropriate patients)
• Long-term cardiovascular risk reduction associated with hypertension
It helps in controlling blood pressure, reducing cardiac workload, and improving cardiovascular health.

Dosage & Administration:
Dosage should be as directed by the physician.
The tablet may be taken with or without food, preferably at the same time each day.
Swallow the tablet whole with water.
Do not discontinue therapy abruptly without medical advice.

Pharmacological Action:

- **Nebivolol is a highly selective β₁-adrenergic receptor antagonist that reduces heart rate and myocardial contractility, thereby decreasing cardiac workload.**
- **It also stimulates endothelial nitric oxide release, producing peripheral vasodilation and improving vascular compliance.**
- **These combined actions effectively lower blood pressure while maintaining favorable cardiovascular hemodynamics.**

Side Effects:
Common side effects include headache, dizziness, fatigue, bradycardia, nausea, diarrhea, constipation, and peripheral edema.
Less commonly, hypotension, sleep disturbances, dyspnea, or hypersensitivity reactions may occur.

Precautions & Warnings:
• Use with caution in patients with bronchial asthma, COPD, diabetes mellitus, peripheral vascular disease, or thyroid disorders.
• Monitor blood pressure and heart rate regularly during treatment.
• Use cautiously in patients with hepatic impairment.
• Do not discontinue therapy abruptly, as sudden withdrawal may worsen cardiovascular conditions.

• Use during pregnancy and lactation only if clearly indicated by the physician.

Drug Interactions:
May interact with calcium channel blockers (e.g., verapamil and diltiazem), antiarrhythmic agents, other antihypertensive medications, digoxin, insulin and oral antidiabetic drugs, anesthetics, and CYP2D6 inhibitors. Clinical monitoring may be required during concomitant therapy.

Storage:
Store below 25°C in a cool, dry place.
Protect from light and moisture.
Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for