Information For the Use of
Nebivolol (5mg) Tablet

Nebiturn 5 Tablet

Nebiturn 5 Tablets

Composition:

Each tablet contains:

- **Nebivolol IP** – 5 mg

Pharmacological Action:

Nebivolol is a **cardioselective β1-adrenergic receptor blocker** with additional vasodilatory activity mediated through nitric oxide.

- **Beta-1 Blockade:** Reduces heart rate, myocardial contractility, and cardiac workload.
- **Vasodilatory Action:** Enhances nitric oxide-mediated vasodilation, reducing peripheral vascular resistance.
- **Antihypertensive Action:** Lowers blood pressure through combined reduction in cardiac workload and peripheral vascular resistance.
- **Cardioprotective Effect:** Reduces myocardial oxygen demand and may help improve cardiovascular efficiency in appropriately selected patients.

Indications:

Nebiturn 5 Tablets is indicated for the management of:

- **Hypertension**
- **Stable chronic heart failure** in appropriately selected patients, as prescribed and monitored by a physician
- Cardiovascular conditions requiring β1-selective beta-blockade, as determined by the physician

---

Dosage and Administration:

- **Recommended Dose:** One tablet **once daily** or as directed by the physician.
- **Administration:** Swallow the tablet whole with water, with or without food.
- Take it at approximately the same time each day.
- Dose may require gradual adjustment according to blood pressure, heart rate, and clinical response.
- **Do not discontinue abruptly**, particularly in patients with ischemic heart disease, unless specifically advised by the physician.

Precautions:

- Monitor **blood pressure and heart rate** regularly.
- Use cautiously in patients with **bradycardia, heart block, sick sinus syndrome, or peripheral vascular disease**.
- Use with caution in patients with **asthma or bronchospastic disease**, despite β1 selectivity.
- Beta-blockers may mask certain symptoms of **hypoglycemia** in diabetic patients.
- Patients with heart failure require careful dose titration and monitoring, particularly during initiation and dose increases.
- Use cautiously in patients with hepatic impairment; dose adjustment may be required.
- **Pregnancy and Lactation:** Use only when clearly indicated and under medical supervision.

Contraindications:

- Known hypersensitivity to nebivolol or any component of the formulation.
- Severe **bradycardia**.
- Second- or third-degree **AV block** without a functioning pacemaker.
- Sick sinus syndrome without a functioning pacemaker.
- **Cardiogenic shock**.
- Decompensated heart failure requiring intensive medical treatment.
- Severe hypotension.
- Severe hepatic impairment where nebivolol use is not appropriate.

Adverse Reactions:

Common:

- Headache
- Dizziness
- Fatigue
- Bradycardia
- Nausea
- Low blood pressure
- Sleep disturbance

Less Common but Serious:

- Significant bradycardia
- Severe hypotension
- Worsening heart failure
- Conduction abnormalities
- Bronchospasm
- Peripheral circulation problems

- Severe hypersensitivity reactions

Drug Interactions:

- **Verapamil and diltiazem** may increase the risk of bradycardia and cardiac conduction abnormalities.
- Other **beta-blockers** or heart-rate-lowering medicines may produce additive effects.
- Other antihypertensive medicines may enhance blood-pressure-lowering effects.
- Certain antiarrhythmic medicines may increase the risk of conduction disturbances.
- CYP2D6 inhibitors may increase nebivolol exposure in some patients.
- Antidiabetic medicines may have their hypoglycemia symptoms masked by beta-blockade.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture and direct sunlight.
- Keep the tablets in their original packaging.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. Nebiturn 5 Tablets should be used under the supervision of a registered medical practitioner. The dosage and duration should be individualized according to the patient's blood pressure, heart rate, cardiovascular status, and clinical response.

Important: Nebivolol should generally not be stopped suddenly, particularly in patients with coronary artery disease, because abrupt withdrawal of beta-blockade can worsen cardiovascular symptoms.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :

Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for