Information For the Use of
Nebivolol (5mg) + Amlodipine (5mg) Tablet

Nemlopine 5/5 Tablet

Nemlopine 5/5 Tablets

Composition:

Each tablet contains:

- **Nebivolol IP** – 5 mg
- **Amlodipine IP** – 5 mg

Pharmacological Action:

Nemlopine 5/5 Tablets combines a cardioselective beta-blocker with a calcium channel blocker to provide complementary antihypertensive activity.

- **Beta-1 Blockade:** Nebivolol selectively blocks β1-adrenergic receptors, reducing heart rate and myocardial workload.
- **Vasodilatory Action:** Nebivolol enhances nitric oxide-mediated vasodilation, helping reduce peripheral vascular resistance.
- **Calcium Channel Blockade:** Amlodipine inhibits calcium influx into vascular smooth muscle, producing prolonged arterial vasodilation.
- **Combined Antihypertensive Action:** The two components reduce vascular resistance and cardiac workload, helping achieve sustained blood pressure control.

Indications:

Nemlopine 5/5 Tablets is indicated for the management of:

- **Hypertension**
- **Hypertension requiring combination therapy**
- **Patients inadequately controlled with monotherapy**, as determined by the physician
- Cardiovascular risk reduction in appropriate hypertensive patients

---

Dosage and Administration:

- **Recommended Dose:** One tablet once daily or as directed by the physician.
- **Administration:** Swallow the tablet whole with water, with or without food.
- Take it at approximately the same time each day.
- Do not discontinue nebivolol abruptly without medical advice.

Precautions:

- Monitor **blood pressure and heart rate** regularly.
- Use cautiously in patients with **bradycardia, conduction abnormalities, peripheral vascular disease, or cardiac dysfunction**.
- Patients with diabetes should be monitored because beta-blockers may mask some symptoms of hypoglycemia.
- Use cautiously in patients with **asthma or bronchospastic disease**.
- Amlodipine may cause peripheral edema; report persistent or severe swelling to the physician.
- Dose selection may require special consideration in patients with hepatic impairment.
- **Pregnancy and Lactation:** Use only when clearly indicated and under medical supervision.

Contraindications:

- Known hypersensitivity to nebivolol, amlodipine, or any component of the formulation.
- Severe bradycardia.
- Significant heart block or sick sinus syndrome without a functioning pacemaker.
- Cardiogenic shock or decompensated heart failure.
- Severe hypotension.
- Severe hepatic impairment where use is not considered appropriate.

Adverse Reactions:

Common:

- Headache
- Dizziness
- Fatigue
- Bradycardia
- Ankle or peripheral edema
- Flushing
- Nausea

Less Common but Serious:

- Excessive hypotension
- Significant bradycardia
- Conduction abnormalities
- Worsening heart failure
- Bronchospasm
- Severe hypersensitivity reactions

Drug Interactions:

- Other **beta-blockers** or heart-rate-lowering medicines may increase the risk of bradycardia.
- **Verapamil and diltiazem** may enhance effects on cardiac conduction and heart rate.
- Other antihypertensive medicines may produce additive blood-pressure-lowering effects.
- CYP3A4 inhibitors may increase amlodipine exposure.
- Certain antidiabetic medicines may have symptoms of hypoglycemia masked by beta-blockade.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture and direct sunlight.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. Nemlopine 5/5 Tablets should be used under the supervision of a registered medical practitioner. The dosage and duration should be individualized according to the patient's blood pressure, heart rate, cardiovascular status, and clinical response.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for