Information For the Use of
Nebivolol (5mg) + Hydrochlorothiazide (12.5mg) Tablet

Nebihydro Tablet

Nebihydro Tablet

Composition:

Each tablet contains:

- **Nebivolol IP** – 5 mg
- **Hydrochlorothiazide IP** – 12.5 mg

Pharmacological Action:

Nebihydro Tablet combines a cardioselective beta-blocker with a thiazide diuretic, providing complementary antihypertensive effects.

- **Beta-1 Blockade:** Nebivolol selectively blocks β1-adrenergic receptors, reducing heart rate and myocardial workload.
- **Vasodilatory Action:** Nebivolol enhances nitric oxide-mediated vasodilation, helping reduce peripheral vascular resistance.
- **Diuretic Action:** Hydrochlorothiazide inhibits sodium and chloride reabsorption in the distal renal tubules, increasing the excretion of sodium and water.
- **Combined Antihypertensive Action:** The combination reduces cardiac workload, circulating fluid volume, and peripheral vascular resistance, resulting in improved blood pressure control.

Indications:

Nebihydro Tablet is indicated for the management of:

- **Hypertension**
- **Hypertension inadequately controlled with monotherapy**
- **Patients requiring combination antihypertensive therapy**, as determined by the physician

---

Dosage and Administration:

- **Recommended Dose:** One tablet once daily or as directed by the physician.
- **Administration:** Swallow the tablet whole with water, with or without food.
- Preferably take the tablet at the same time each day.
- Since hydrochlorothiazide increases urine production, morning administration may be preferable.
- Do not discontinue nebivolol abruptly without medical advice.

Precautions:

- Monitor **blood pressure and heart rate** regularly.
- Periodic monitoring of **renal function and serum electrolytes** may be required.
- Hydrochlorothiazide may alter potassium, sodium, glucose, uric acid, and calcium levels.
- Use cautiously in patients with **diabetes mellitus, gout, renal impairment, or electrolyte disturbances**.
- Nebivolol may mask certain symptoms of hypoglycemia.
- Use cautiously in patients with **asthma or bronchospastic disease**.
- Maintain adequate fluid intake, particularly during excessive sweating, vomiting, or diarrhea.
- **Pregnancy and Lactation:** Use only when clearly indicated and under medical supervision.

Contraindications:

- Known hypersensitivity to nebivolol, hydrochlorothiazide, sulfonamide-derived medicines, or any component of the formulation.
- Severe bradycardia.
- Significant heart block or sick sinus syndrome without a functioning pacemaker.
- Cardiogenic shock or decompensated heart failure.
- Severe hypotension.
- **Anuria** or severe renal impairment where hydrochlorothiazide is inappropriate.
- Severe hepatic impairment requiring specialist assessment.

Adverse Reactions:

Common:

- Dizziness
- Fatigue
- Headache
- Bradycardia
- Increased urination
- Nausea
- Electrolyte disturbances

Less Common but Serious:

- Excessive hypotension
- Significant bradycardia
- Bronchospasm
- Electrolyte imbalance
- Increased uric acid levels or gout
- Worsening renal function

- Severe hypersensitivity reactions

Drug Interactions:

- Other **beta-blockers** or heart-rate-lowering medicines may increase the risk of bradycardia.
- **Verapamil and diltiazem** may enhance effects on cardiac conduction and heart rate.
- Other antihypertensive medicines may produce additive blood-pressure-lowering effects.
- **NSAIDs** may reduce the antihypertensive effect and may increase the risk of renal impairment.
- **Lithium** levels may increase with thiazide diuretics.
- Antidiabetic medicines may require monitoring because hydrochlorothiazide can affect glucose control.
- Medicines affecting potassium levels may increase the risk of electrolyte disturbances.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture and direct sunlight.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. Nebihydro Tablet should be used under the supervision of a registered medical practitioner. The dosage and duration of treatment should be individualized according to the patient's blood pressure, heart rate, renal function, electrolyte status, and clinical response.

For Nebihydro, the main safety distinction from Nemlopine is the added diuretic effect, so compare Nebihydro with Nemlopine by benefits, side effects, and monitoring needs before choosing between them.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :

Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for