Information For the Use of
Nepafenac (0.1% w/v) Eye Drops

Nepastep Eye Drops

Brand Name: Nepastep Eye Drops
Dosage Form: Eye Drops
Composition:
Each ml contains:
• Nepafenac IP 0.1% w/v
• Sterile ophthalmic vehicle q.s.

Description:
Nepastep Eye Drops contains Nepafenac, a non-steroidal anti-inflammatory drug (NSAID) formulated for ophthalmic use. It is indicated for the prevention and treatment of pain and inflammation associated with ocular surgical procedures, particularly cataract surgery. It helps in reducing ocular inflammation, redness, swelling, and postoperative discomfort by inhibiting prostaglandin synthesis within ocular tissues.

Indications:
Indicated for the treatment and prevention of:
• Post-operative ocular pain and inflammation
• Inflammation associated with cataract surgery
• Ocular discomfort following ophthalmic procedures
• Sterile inflammatory conditions of the eye as advised by the physician
It helps in reducing pain, redness, swelling, and inflammation of the eye.

Dosage & Administration:
Instill one drop into the affected eye(s) as directed by the physician.
Wash hands thoroughly before administration.
Avoid touching the dropper tip to any surface to prevent contamination.
If using multiple ophthalmic medications, maintain an interval of at least 5–10 minutes between applications.

Pharmacological Action:

- **Nepafenac is a prodrug that rapidly penetrates ocular tissues and is converted to amfenac, its active metabolite.**
- **Amfenac inhibits cyclooxygenase (COX) enzymes, thereby reducing prostaglandin synthesis responsible for ocular pain and inflammation.**

These actions help in effective control of postoperative inflammation and ocular discomfort.

Side Effects:
Common side effects include transient burning or stinging sensation, eye irritation, blurred vision, foreign body sensation, and increased lacrimation.
Rarely, corneal disorders, delayed healing, or hypersensitivity reactions may occur.

Precautions & Warnings:
• For ophthalmic use only
• Use with caution in patients with known hypersensitivity to NSAIDs
• Prolonged use may increase the risk of corneal adverse events in susceptible patients

• Monitor patients carefully following ophthalmic surgery
• Avoid contamination of the dropper tip
• Remove contact lenses before administration and reinsert as advised by the physician

---

Drug Interactions:
May be administered with other ophthalmic medications under medical supervision.
Concurrent use with topical corticosteroids may delay wound healing.

Storage:
Store below 25°C in a cool, dry place.
Protect from light.
Do not freeze.
Keep the bottle tightly closed after use.
Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for