Critsila™

Nestomin-2.5
Injection

Neostigmine 2.5mg





Rapid & Reliable Action
Quickly reverses non-depolarizing neuromuscular blockade

Proven Reversal Agent
Gold standard for reversal of neuromuscular blocking agents

Restores Neuromuscular Function
Enhances acetylcholine activity at the neuromuscular junction

Predictable & Controllable
Short duration of action allows better titration & control

Well Tolerated
Extensively used with established safety profile in clinical practice

Broad Clinical Utility
Useful in anesthesia, ICU & various clinical settings

Indications

- Reversal of Non-depolarizing Neuromuscular Blockade
- Postoperative Recovery of Muscle Function
- ICU & Critical Care Settings
- Paralytic Ileus & Bowel Dysfunction
- Postoperative Urinary Retention

References

1. Savarese JJ, Ali HH, Basta SJ. Neostigmine. [Updated 2023 Aug 14]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: https://www.ncbi.nlm.nih.gov/books/NBK482354/
2. McEvoy GK. AHFS Drug Information 2024. American Society of Health-System Pharmacists; 2024.
3. Miller RD, Eriksson LI, Fleisher LA, Wiener-Kronish JP, Young WL. Miller's Anesthesia. 9th ed. Philadelphia: Elsevier; 2020.