Information For the Use of
Olmesartan medoxomil 20 mg + amlodipine 5 mg + hydrochlorothiazide 12.5 mg tablet

Olmeslide AH 20 Tablet

Brand Name: Olmeslide AH 20 Tablet
Dosage Form: Tablet
Composition:
Each film-coated tablet contains:
• Olmesartan Medoxomil IP 20 mg
• Amlodipine Besylate IP equivalent to Amlodipine 5 mg
• Hydrochlorothiazide IP 12.5 mg

Description:
Olmeslide AH 20 Tablet is a fixed-dose combination of Olmesartan Medoxomil, an angiotensin II receptor blocker (ARB); Amlodipine, a calcium channel blocker; and Hydrochlorothiazide, a thiazide diuretic. It is indicated for the management of hypertension in patients requiring combination therapy. It helps in controlling blood pressure through complementary mechanisms, thereby reducing cardiovascular risk and improving vascular health.

Indications:
Indicated for the treatment of:
• Essential hypertension
• Hypertension inadequately controlled with dual therapy
It helps in lowering elevated blood pressure and reducing the risk of cardiovascular complications such as stroke and myocardial infarction.

Dosage & Administration:
Dosage should be as directed by the physician.
The tablet may be taken with or without food, preferably at the same time each day.
Regular monitoring of blood pressure, renal function, and electrolytes is recommended during therapy.

Pharmacological Action:

- **Olmesartan Medoxomil blocks angiotensin II receptors, resulting in vasodilation and reduced aldosterone secretion.**
- **Amlodipine inhibits calcium ion influx into vascular smooth muscle, producing peripheral vasodilation.**
- **Hydrochlorothiazide promotes sodium and water excretion by inhibiting sodium reabsorption in the distal renal tubules.**

Together, these actions provide effective and sustained blood pressure control.

Side Effects:
Common side effects include dizziness, headache, fatigue, ankle edema, increased urination, and electrolyte imbalance.
Rarely, hypotension, renal impairment, or hypersensitivity reactions may occur.

Precautions & Warnings:

• Use with caution in patients with renal or hepatic impairment
• Monitor serum electrolytes and renal function periodically
• Risk of hypotension may increase in volume-depleted patients
• Use cautiously in diabetic patients and those with gout
• Not recommended during pregnancy

Drug Interactions:
May interact with other antihypertensive agents, lithium, NSAIDs, corticosteroids, and potassium supplements.
Concomitant use with alcohol may enhance hypotensive effects.

Storage:
Store below 25°C in a cool, dry place.
Protect from moisture and light.
Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for