Information For the Use of
Choline Salicylate (8.7% w/w) + Lidocaine (2.0% w/w) + Benzalkonium Chloride (0.01% w/w) Gel

Ovastyle Ulcer Gel

Brand Name: Ovastyle Ulcer Gel
Dosage Form: Oral Topical Gel
Composition:
Each gram contains:
• Choline Salicylate IP 8.7% w/w
• Lidocaine Hydrochloride IP 2.0% w/w
• Benzalkonium Chloride IP 0.01% w/w
• Gel base q.s.

Description:
Ovastyle Ulcer Gel is a combination oral mucosal analgesic and anti-inflammatory gel formulated for the management of painful oral ulcers and related conditions. It provides rapid relief from pain and inflammation while promoting patient comfort during eating and speaking. It helps in soothing irritated oral mucosa and reducing discomfort associated with ulcerative lesions.

Indications:
Indicated for the treatment of:
• Recurrent aphthous ulcers
• Oral mucositis
• Gingival inflammation
• Minor oral lesions and irritation
It helps in relieving pain, reducing inflammation, and protecting the oral mucosal surface.

Dosage & Administration:
Apply a small amount of gel to the affected area 3–4 times daily or as directed by the physician.
Avoid eating or drinking for at least 30 minutes after application for optimal effect.
Use with a clean fingertip or applicator.

Pharmacological Action:

- **Choline Salicylate acts as a local anti-inflammatory agent by inhibiting prostaglandin synthesis, thereby reducing pain and swelling.**
- **Lidocaine is a local anesthetic that blocks sodium channels, providing rapid pain relief.**
- **Benzalkonium Chloride acts as an antiseptic, reducing microbial load and preventing secondary infection.**

Together, these components provide analgesic, anti-inflammatory, and antiseptic effects in oral lesions.

Side Effects:
Common side effects include mild burning sensation, local irritation, or numbness at the application site.
Rarely, hypersensitivity reactions may occur.

Precautions & Warnings:
• For oral topical use only
• Avoid swallowing large quantities

- **Use with caution in patients with salicylate or local anesthetic sensitivity**
- **Discontinue use if irritation persists or worsens**
- **Keep out of reach of children**

Drug Interactions:
No significant systemic interactions reported due to minimal absorption.
However, caution is advised with other salicylate-containing products.

Storage:
Store below 25°C in a cool, dry place.
Do not freeze.
Keep tube tightly closed after use.
Protect from direct sunlight.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for