Information For the Use of
Caffeine (30mg) + Diphenhydramine (25mg) + Paracetamol (500mg) + Phenylephrine (5mg) Tablets

Coldstave Cold N Flu Tablet

COLDSTAVE COLD N FLU TABLET

Composition

Each film-coated tablet contains:

Paracetamol I.P. 500 mg
Phenylephrine Hydrochloride I.P. 5 mg
Caffeine I.P. 30 mg
Diphenhydramine Hydrochloride I.P. 25 mg
Excipients … q.s.

Pharmacological Action

Paracetamol is an analgesic and antipyretic that provides relief from fever, headache and mild-to-moderate body aches associated with common cold and influenza.

Phenylephrine Hydrochloride is a sympathomimetic nasal decongestant. It produces vasoconstriction of the nasal mucosal blood vessels, thereby reducing nasal congestion and stuffiness.

Diphenhydramine Hydrochloride is a first-generation H1-antihistamine that helps relieve histamine-mediated symptoms such as sneezing, rhinorrhoea and watery eyes. It may also cause drowsiness.

Caffeine is a central nervous system stimulant and may enhance the analgesic effect of paracetamol while helping to counteract some antihistamine-associated drowsiness.

Together, the formulation provides **analgesic, antipyretic, antihistaminic and nasal-decongestant activity** for symptomatic relief of common cold and influenza-associated symptoms.

Indications

Coldstave Cold N Flu Tablet is indicated for symptomatic relief of:

- Common cold and influenza symptoms
- Fever and headache
- Body ache associated with cold and flu
- Nasal congestion and blocked nose
- Sneezing and runny nose

- Watery eyes and other allergy-associated cold symptoms

It is intended for **short-term symptomatic treatment** and does not treat the underlying viral infection.

Dosage and Administration

Adults and children above 12 years:
1 tablet every **6–8 hours as required**, or as directed by the physician.

Maximum: Do not exceed **4 tablets in 24 hours** unless specifically advised by a physician.

Swallow the tablet whole with water. Do not chew or crush.

Use the lowest effective dose for the shortest necessary duration. If symptoms persist, worsen, or are accompanied by significant fever, medical evaluation is recommended.

Precautions

- Use cautiously in patients with **hypertension, cardiovascular disease or hyperthyroidism** because of the phenylephrine component.
- Use cautiously in patients with diabetes, glaucoma, urinary retention or prostate enlargement.
- Diphenhydramine may cause **drowsiness, dizziness and impaired alertness**; avoid driving or operating machinery if affected.
- Avoid alcohol and other medicines that cause CNS depression.
- Do not take with other medicines containing **paracetamol**, as excessive total paracetamol intake may cause severe liver injury.
- Caffeine may cause restlessness, insomnia or palpitations in susceptible individuals.
- Patients with hepatic or renal impairment should use the medicine only under medical advice.
- Do not use continuously for prolonged periods without medical advice.

Contraindications

Coldstave Cold N Flu Tablet is contraindicated in patients with:

- Hypersensitivity to paracetamol, phenylephrine, diphenhydramine, caffeine or any component of the formulation
- Severe hypertension or significant cardiovascular disease
- Closed-angle glaucoma
- Severe hepatic impairment
- Urinary retention associated with significant bladder outlet obstruction
- Concomitant use of **monoamine oxidase inhibitors (MAOIs)** or use within the preceding 14 days
- Concomitant use of other sympathomimetic decongestants

Use in children below 12 years should be based on medical advice.

Adverse Reactions

Common/possible:

- Drowsiness
- Dizziness
- Dry mouth
- Nausea or gastric discomfort
- Headache
- Blurred vision
- Restlessness or nervousness
- Insomnia
- Palpitations

Rare/serious:

- Significant increase in blood pressure or heart rate
- Urinary retention
- Severe hypersensitivity reactions
- Severe skin reactions
- Liver injury following paracetamol overdose or excessive use

Drug Interactions

MAO inhibitors: Concomitant use with phenylephrine may cause a potentially dangerous increase in blood pressure and is contraindicated.

Other sympathomimetics/decongestants: May increase cardiovascular adverse effects.

Beta-blockers and antihypertensive medicines: Phenylephrine may reduce the effectiveness of some antihypertensive agents.

Sedatives, hypnotics and CNS depressants: May enhance the sedative effects of diphenhydramine.

Anticholinergic medicines: May increase anticholinergic adverse effects such as dry mouth, blurred vision and urinary retention.

Paracetamol-containing medicines: Concurrent use increases the risk of accidental paracetamol overdose and hepatotoxicity.

Warfarin: Prolonged regular use of paracetamol may enhance anticoagulant effect in some patients.

Caffeine: Concomitant intake of other caffeine-containing products may increase nervousness, insomnia, tremor or palpitations. Phenylephrine and caffeine may both increase blood pressure and heart rate.

Pregnancy and Lactation

Use during pregnancy or lactation should be undertaken only after consultation with a healthcare professional, particularly because the formulation contains a systemic decongestant and first-generation antihistamine.

Storage

Store **below 30°C**, in a cool, dry place, protected from light and moisture.

Keep the medicine in its original packaging and out of reach of children.

Note

For use under medical supervision.

Do not exceed the recommended dose. Avoid taking other cold, flu or pain-relief preparations containing **paracetamol, phenylephrine, diphenhydramine or caffeine** without professional advice. If symptoms persist or worsen, consult a physician.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for