Pirfound 200 Tablet

PIRFOUND 200 TABLETS

Composition

Each film-coated tablet contains:

Pirfenidone 200 mg
Excipients … q.s.

Pharmacological Action

Pirfenidone is an oral antifibrotic agent with anti-inflammatory and antifibrotic properties.

It is believed to reduce the production of pro-fibrotic mediators and cytokines, including transforming growth factor-β (TGF-β), thereby reducing fibroblast proliferation, collagen synthesis and accumulation of extracellular matrix.

Pirfenidone may help **slow the decline in lung function** associated with pulmonary fibrosis.

Indications

Pirfound 200 Tablets are indicated for the treatment of:

- **Idiopathic Pulmonary Fibrosis (IPF)** in adults.

Pirfenidone is used to slow disease progression and decline in pulmonary function in patients with IPF.

Dosage and Administration

Adults:

Pirfenidone therapy is generally initiated with a gradual dose escalation:

- **Days 1–7:** 200 mg three times daily
- **Days 8–14:** 400 mg three times daily
- **Day 15 onward:** 600 mg three times daily

The usual maintenance dosage is therefore **600 mg three times daily (1,800 mg/day)**, as tolerated and prescribed by the physician.

Pirfound 200 Tablets should be taken **with food** to improve gastrointestinal tolerability.

Tablets should be swallowed whole with water.

If treatment is interrupted for a prolonged period, re-titration may be required according to the prescribing information and physician's advice.

Precautions

- Liver-function tests should be performed before initiating treatment and monitored periodically during therapy.
- Use cautiously in patients with pre-existing hepatic impairment.
- Monitor for symptoms or laboratory evidence of **drug-induced liver injury**.
- Patients should be advised to minimize exposure to direct sunlight and use protective clothing and sunscreen because **photosensitivity reactions** may occur.
- Gastrointestinal adverse effects such as nausea, dyspepsia and diarrhoea may occur, particularly during dose escalation.
- Smoking may reduce exposure to pirfenidone and may decrease its therapeutic effect; patients should be advised not to smoke.
- Use cautiously in patients with renal impairment.
- Report severe rash, blistering, marked photosensitivity, jaundice, dark urine or unusual fatigue promptly to the physician.

Contraindications

Pirfound 200 Tablets are contraindicated in:

- Known hypersensitivity to **pirfenidone** or any component of the formulation.
- Concomitant use with **fluvoxamine**, unless specifically managed according to the approved prescribing information.
- Severe hepatic impairment or end-stage liver disease where pirfenidone is not appropriate.
- Severe renal impairment requiring dialysis, where use is generally not recommended.

Contraindications and dose restrictions should be confirmed against the applicable approved product information.

Adverse Reactions

Common:

- Nausea
- Diarrhoea
- Dyspepsia
- Vomiting
- Abdominal discomfort
- Fatigue
- Headache
- Dizziness
- Decreased appetite

- Weight loss

Skin-related:

- Photosensitivity reaction
- Rash
- Pruritus

Laboratory abnormalities:

- Increased ALT/AST
- Other liver-function abnormalities

Serious/rare:

- Clinically significant liver injury
- Severe photosensitivity or serious skin reactions

Dose reduction or temporary interruption may be required if significant adverse reactions occur.

Drug Interactions

Fluvoxamine:
Concomitant use can substantially increase pirfenidone exposure and should generally be avoided.

CYP1A2 inhibitors:
Other strong inhibitors of CYP1A2 may increase pirfenidone exposure and may require dose adjustment according to prescribing information.

CYP1A2 inducers:
Strong CYP1A2 inducers, including smoking, can reduce pirfenidone exposure and potentially reduce its effectiveness.

Patients should inform their physician about all prescription medicines, OTC medicines, herbal products and supplements being used.

Pregnancy and Lactation

Pirfenidone should generally **not be used during pregnancy unless clearly necessary and prescribed by a specialist**.

Women of reproductive potential should discuss pregnancy planning with their physician before starting treatment.

It is not established whether pirfenidone or its metabolites are present in human milk. Breastfeeding should be discussed with the treating physician before treatment.

Storage

Store in a **cool, dry place**, protected from excessive heat, moisture and direct sunlight.

Keep the tablets in their original packaging and **out of reach of children**.

Note

For use under medical supervision.

Pirfound 200 Tablets should be used only for the prescribed indication. Regular monitoring of **liver function, tolerability and clinical status** is recommended during treatment.

The exact dose, dose modifications and treatment duration should be determined by the treating physician according to the patient's clinical condition and the approved product prescribing information.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for