Information For the Use of
Prazosin (2.5mg) Sustained Release Tablet

Prazotyle XL 2.5 Tablet

Brand Name: Prazotyle XL 2.5 Tablet
Dosage Form: Sustained Release (SR) Tablet
Composition:
Each sustained release tablet contains:
• Prazosin Hydrochloride IP equivalent to Prazosin 2.5 mg (Sustained Release)

Description:
Prazotyle XL 2.5 Tablet contains Prazosin, a selective α₁-adrenergic receptor antagonist formulated as a sustained release tablet. It is indicated for the management of essential hypertension by reducing peripheral vascular resistance through selective α₁-receptor blockade. The sustained release formulation ensures prolonged drug release, providing sustained blood pressure control and improved patient compliance.

Indications:
Indicated for the treatment of:
• Essential hypertension
• Adjunctive therapy in resistant hypertension
• Benign prostatic hyperplasia (BPH) with associated lower urinary tract symptoms, where clinically appropriate
It helps in lowering blood pressure, reducing cardiac workload, improving peripheral circulation, and relieving urinary symptoms associated with benign prostatic hyperplasia.

Dosage & Administration:
Dosage should be as directed by the physician.
The sustained release tablet should be swallowed whole with water and should not be crushed, chewed, or broken.
It may be taken with or without food, preferably at the same time each day.
The first dose is preferably administered at bedtime to reduce the risk of first-dose hypotension.

Pharmacological Action:

- **Prazosin selectively blocks post-synaptic α₁-adrenergic receptors in vascular smooth muscle, producing arterial and venous vasodilation.**
- **It decreases peripheral vascular resistance, lowers systemic blood pressure, and reduces cardiac afterload without significantly affecting cardiac output.**
- **In patients with benign prostatic hyperplasia, Prazosin relaxes smooth muscle in the bladder neck and prostate, thereby improving urinary flow and reducing lower urinary tract symptoms.**

Side Effects:
Common side effects include dizziness, headache, fatigue, drowsiness, weakness, nausea, palpitations, and postural hypotension.
Occasionally, syncope, edema, blurred vision, nasal congestion, or hypersensitivity reactions may occur.

Precautions & Warnings:

• Initiate therapy cautiously because of the risk of first-dose hypotension and syncope.
• Monitor blood pressure regularly during treatment.
• Advise patients to rise slowly from sitting or lying positions to minimize orthostatic hypotension.
• Use with caution in patients with hepatic impairment, severe cardiovascular disease, or renal impairment.
• Use during pregnancy and lactation only if the potential benefit justifies the potential risk, as determined by the physician.

Drug Interactions:
May interact with other antihypertensive agents, diuretics, beta-blockers, phosphodiesterase-5 (PDE-5) inhibitors (e.g., sildenafil, tadalafil), and alcohol, increasing the risk of hypotension. Clinical monitoring is recommended during concomitant therapy.

Storage:
Store below 25°C in a cool, dry place.
Protect from light and moisture.
Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for