Information For the Use of
Prazosin (5 mg) Sustained Release Tablet

Prazotyle XL 5 Tablet

Brand Name: Prazotyle XL 5 Tablet
Dosage Form: Sustained Release (SR) Tablet
Composition:
Each sustained release tablet contains:
• Prazosin Hydrochloride IP equivalent to Prazosin 5 mg (Sustained Release)

Description:
Prazotyle XL 5 Tablet contains Prazosin, a selective α₁-adrenergic receptor antagonist formulated as a sustained release tablet. It is indicated for the management of essential hypertension by producing peripheral vasodilation and reducing systemic vascular resistance. The sustained release formulation provides prolonged therapeutic action, ensuring consistent blood pressure control throughout the day while improving patient compliance.

Indications:
Indicated for the treatment of:
• Essential hypertension
• Adjunctive therapy in resistant hypertension
• Benign prostatic hyperplasia (BPH) with associated lower urinary tract symptoms, where clinically appropriate
It helps in lowering blood pressure, reducing peripheral vascular resistance, improving urinary flow, and relieving symptoms associated with benign prostatic hyperplasia.

Dosage & Administration:
Dosage should be as directed by the physician.
The sustained release tablet should be swallowed whole with water and should not be crushed, chewed, or broken.
It may be taken with or without food, preferably at the same time each day.
The initial dose should preferably be administered at bedtime to minimize the risk of first-dose hypotension.

Pharmacological Action:

- **Prazosin selectively blocks post-synaptic α₁-adrenergic receptors in vascular smooth muscle, producing arterial and venous vasodilation.**
- **It reduces peripheral vascular resistance, decreases blood pressure, and lowers cardiac afterload with minimal effect on cardiac output.**
- **In patients with benign prostatic hyperplasia, it relaxes smooth muscle of the prostate and bladder neck, thereby improving urinary flow and reducing lower urinary tract symptoms.**

Side Effects:
Common side effects include dizziness, headache, fatigue, drowsiness, weakness, nausea, palpitations, postural hypotension, and nasal congestion.
Occasionally, syncope, edema, blurred vision, or hypersensitivity reactions may occur.

Precautions & Warnings:
• Initiate therapy cautiously because of the risk of first-dose hypotension and syncope.

• Monitor blood pressure regularly during treatment.
• Advise patients to rise slowly from sitting or lying positions to reduce orthostatic hypotension.
• Use with caution in patients with hepatic impairment, renal impairment, or severe cardiovascular disease.
• Use during pregnancy and lactation only if clearly indicated by the physician.

Drug Interactions:
May interact with other antihypertensive agents, diuretics, beta-blockers, phosphodiesterase-5 (PDE-5) inhibitors (e.g., sildenafil, tadalafil), and alcohol, increasing the risk of hypotension. Clinical monitoring is recommended during concomitant therapy.

Storage:
Store below 25°C in a cool, dry place.
Protect from light and moisture.
Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for