Information For the Use of
Pyrimethamine (37.5mg) + Sulphadoxine (750mg) Tablet

Pyrethon S Tablet

Pyrethon S Tablet

Composition:

Each tablet contains:

- **Pyrimethamine IP** – 37.5 mg
- **Sulphadoxine IP** – 750 mg

Pharmacological Action:

Pyrethon S Tablet is a fixed-dose combination of **Pyrimethamine** and **Sulphadoxine**, two antifolate agents with synergistic antimalarial activity. The combination acts by sequentially inhibiting folate synthesis in the malaria parasite, thereby preventing nucleic acid synthesis and parasite replication.

- **Pyrimethamine:** Selectively inhibits parasitic **dihydrofolate reductase (DHFR)**, interfering with folic acid metabolism required for DNA synthesis.
- **Sulphadoxine:** Inhibits **dihydropteroate synthase (DHPS)**, blocking the early step of folate synthesis in the parasite.
- **Synergistic Action:** Sequential blockade of folate metabolism enhances antimalarial efficacy and delays parasite growth.
- **Long Duration of Action:** The long half-life of both components provides sustained therapeutic activity.

Indications:

Pyrethon S Tablet is indicated for the treatment of susceptible **Plasmodium falciparum** malaria and other protozoal infections where appropriate.

It is used in the management of:

- **Uncomplicated Plasmodium falciparum malaria** (in combination with recommended antimalarial agents as per current treatment guidelines).
- **Mixed malarial infections** where indicated.
- **Toxoplasmosis** (in combination with other appropriate antimicrobial agents, as advised by the physician).

Dosage and Administration:

- **Recommended Dose:** As directed by the physician and according to national malaria treatment guidelines.
- **Administration:** Swallow the tablet whole with water, preferably after food to reduce gastrointestinal discomfort.
- Adequate hydration should be maintained during therapy.
- Complete the full prescribed course even if symptoms improve earlier.

Precautions:

- Use with caution in patients with **hepatic or renal impairment**.
- Caution is advised in patients with **folate deficiency**, malnutrition, or hematological disorders.
- Periodic blood counts may be required during prolonged therapy.
- Use cautiously in patients with **glucose-6-phosphate dehydrogenase (G6PD) deficiency**.
- Pregnancy and lactation: Use only if clearly needed and under medical supervision.

Contraindications:

- Known hypersensitivity to **Pyrimethamine**, **Sulphonamides**, or any component of the formulation.
- Severe hepatic or renal impairment.
- Patients with megaloblastic anemia due to folate deficiency.
- Infants below the recommended age unless specifically prescribed.

Adverse Reactions:

Common:

- Nausea
- Vomiting
- Loss of appetite
- Headache
- Abdominal discomfort

Less Common but Serious:

- Skin rash
- Stevens–Johnson syndrome (rare)
- Toxic epidermal necrolysis (rare)
- Blood dyscrasias (anemia, leukopenia, thrombocytopenia)

- Allergic reactions

Patients should seek immediate medical attention if severe skin reactions, persistent fever, sore throat, unusual bleeding, or allergic symptoms occur.

Drug Interactions:

- Concomitant use with other **folate antagonists** (e.g., methotrexate, trimethoprim) may increase the risk of bone marrow suppression.
- May potentiate the effects of **oral anticoagulants**.
- Caution with drugs causing bone marrow toxicity.
- Folinic acid may be required in selected patients receiving prolonged therapy.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from light and moisture.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. Pyrethon S Tablet should be used only under the supervision of a registered medical practitioner and in accordance with current national malaria treatment guidelines. The prescribing physician should consider local antimalarial resistance patterns and individual patient factors before initiating therapy.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for