Information For the Use of
Quinine (300mg) Injection

Quinshell 300 Injection

QUINSHELL 300 INJECTION

Quinine 300 mg Injection

COMPOSITION

Each ampoule/vial contains:

Quinine 300 mg
Suitable Sterile Vehicle q.s.

The exact salt form and concentration should be confirmed against the approved product specification before final printing.

PHARMACOLOGICAL ACTION

Quinine is an antimalarial agent derived from the bark of the cinchona tree. It acts against the **erythrocytic stages of Plasmodium parasites**, interfering with parasite haemoglobin metabolism and contributing to parasite death.

Quinine has activity against *Plasmodium falciparum* and other malaria parasites, although resistance and safety considerations limit its use compared with newer antimalarial therapies.

INDICATIONS

Quinshell 300 Injection may be used under specialist supervision for:

- **Severe or complicated malaria**, particularly when parenteral therapy is required
- Severe *Plasmodium falciparum* malaria when recommended alternatives are unavailable or unsuitable
- Malaria in patients unable to tolerate or receive oral antimalarial therapy

Treatment should follow current national malaria-treatment guidelines and local resistance patterns.

DOSAGE AND ADMINISTRATION

Quinine injection must be administered by a qualified healthcare professional in a hospital or appropriate clinical setting.

For severe malaria, the dose is **weight-based** and depends on the quinine salt/formulation, patient's age, previous antimalarial therapy and local treatment guidelines.

Where quinine dihydrochloride is used for severe malaria, commonly referenced regimens include an initial **20 mg/kg salt (maximum 1.4 g) IV loading dose**, followed by **10 mg/kg every 8 hours**, with appropriate adjustment according to clinical response and toxicity.

- Administer by **slow intravenous infusion after appropriate dilution**.
- **Do not administer by rapid IV injection.**
- Monitor blood glucose, ECG where indicated, renal function and clinical response.
- Oral antimalarial therapy should replace parenteral treatment as soon as the patient can tolerate it, followed by completion of an appropriate full antimalarial regimen.

The exact dose and concentration must be calculated from the approved formulation and current national malaria guidelines.

PRECAUTIONS

- Quinine may cause **hypoglycaemia**, particularly in severe malaria and during pregnancy; monitor blood glucose regularly.
- Monitor for **QT prolongation and cardiac arrhythmias**, especially in patients with cardiac disease or electrolyte abnormalities.
- Use cautiously in patients with hepatic or renal impairment.
- Monitor for tinnitus, hearing disturbance, headache, dizziness and visual symptoms.
- Quinine may cause thrombocytopenia and haemolytic reactions in susceptible patients.
- Avoid dosing errors because quinine has a relatively narrow therapeutic range.
- Patients receiving repeated doses should be monitored for cinchonism and other toxicity.
- Adequate treatment of hypoglycaemia, anaemia, renal impairment and other complications of severe malaria is essential.

CONTRAINDICATIONS

- Hypersensitivity to quinine or quinidine
- Previous quinine-induced thrombocytopenia or other serious quinine-related immune reaction

- Significant cardiac conduction abnormalities or clinically important QT prolongation, unless specialist assessment determines otherwise
- Myasthenia gravis, where quinine may worsen neuromuscular weakness
- Optic neuritis or significant previous quinine-associated visual toxicity

---

ADVERSE REACTIONS

Common

- Headache
- Nausea and vomiting
- Abdominal discomfort
- Dizziness
- Tinnitus
- Hearing disturbance
- Blurred vision
- Sweating
- Weakness

These symptoms are collectively associated with **cinchonism**.

Serious / Rare

- Hypoglycaemia
- QT prolongation and cardiac arrhythmias
- Hypotension
- Thrombocytopenia
- Haemolytic anaemia
- Severe hypersensitivity reactions
- Bronchospasm
- Acute kidney injury
- Severe visual or auditory disturbances

DRUG INTERACTIONS

Quinine may interact with medicines that affect cardiac conduction or its metabolism.

Important interactions include:

- Quinidine and other antiarrhythmic medicines
- Medicines that prolong the QT interval

- Warfarin and other anticoagulants
- Digoxin
- Certain macrolide antibiotics
- Rifampicin and other enzyme-inducing medicines
- Cimetidine
- Antidiabetic medicines, because of increased risk of hypoglycaemia

Concomitant medicines should be reviewed before administration.

USE IN PREGNANCY AND LACTATION

Pregnancy:
Severe malaria is a medical emergency and must be treated promptly. Quinine may be used during pregnancy when clinically indicated, particularly when recommended by current malaria-treatment guidelines. Maternal blood glucose should be monitored closely because quinine can cause hypoglycaemia.

Lactation:
Small amounts of quinine may pass into breast milk. Use during breastfeeding should be based on clinical necessity and physician assessment.

STORAGE

Store according to the approved product label, generally protected from light and excessive heat.

Do not freeze.

Keep out of reach of children.

NOTE

Quinshell 300 Injection is intended for **hospital/clinical use under qualified medical supervision**.

Severe malaria requires urgent treatment and supportive management. The **exact quinine salt, concentration, dose and infusion regimen must be verified from the approved product specification and current Indian malaria-treatment guidelines before finalizing printed literature.**

For professional use only.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for