Information For the Use of
Ramipril (5mg) + Hydrochlorothiazide (12.5mg) Tablet

Rampolite H Tablet

Rampolite H Tablet

Composition:

Each tablet contains:

- **Ramipril** – 5 mg
- **Hydrochlorothiazide** – 12.5 mg

Pharmacological Action:

Rampolite H combines an **ACE inhibitor** with a **thiazide diuretic** to provide complementary antihypertensive effects.

- **ACE Inhibition:** Ramipril inhibits angiotensin-converting enzyme, reducing formation of angiotensin II and aldosterone, resulting in vasodilation and reduced sodium and water retention.
- **Diuretic Action:** Hydrochlorothiazide increases renal excretion of sodium and water, reducing plasma volume and peripheral vascular resistance.
- **Complementary Action:** The combination provides enhanced blood-pressure reduction compared with either component alone in appropriately selected patients.
- **Cardiovascular Effect:** Reduced vascular resistance and fluid volume decrease cardiac workload.

Indications:

Rampolite H Tablet is indicated for the management of:

- **Hypertension**
- **Essential Hypertension**
- Hypertension inadequately controlled with monotherapy, when combination therapy is appropriate
- Patients requiring an ACE inhibitor with a diuretic for blood-pressure control

---

Dosage and Administration:

- **Recommended Dose:** One tablet once daily or as directed by the physician.
- Preferably take the tablet at the **same time each day**, usually in the morning because of the diuretic effect.
- Swallow the tablet whole with water.

- It may be taken with or without food.
- Blood pressure, renal function, and electrolytes should be monitored periodically.
- Dose adjustments should be made by the physician according to clinical response.

Precautions:

- Monitor **blood pressure, renal function, serum potassium, and sodium** during treatment.
- Ramipril may cause **hyperkalemia**, while hydrochlorothiazide may cause electrolyte depletion.
- Use cautiously in patients with **renal impairment, hepatic impairment, dehydration, or electrolyte abnormalities**.
- ACE inhibitors may cause a persistent **dry cough** and, rarely, angioedema.
- Patients with diabetes should have blood glucose monitored because hydrochlorothiazide may affect glucose control.
- Use cautiously in patients with a history of **gout**, as hydrochlorothiazide may increase uric acid levels.
- Excessive fluid loss from vomiting, diarrhea, or heavy sweating may increase the risk of hypotension and renal impairment.
- **Pregnancy:** ACE inhibitors can cause fetal harm and should not be used during pregnancy.
- **Lactation:** Consult a physician before use.

Contraindications:

- Known hypersensitivity to ramipril, hydrochlorothiazide, other ACE inhibitors, sulfonamide-derived diuretics, or any component of the formulation.
- **History of ACE-inhibitor-associated angioedema**.
- **Pregnancy**.
- Severe renal impairment or anuria where the combination is not appropriate.
- Significant electrolyte disturbances that have not been corrected.
- Concomitant use with **aliskiren in patients with diabetes or significant renal impairment**.
- Concomitant use with certain **neprilysin inhibitors** requires appropriate separation and specialist assessment.

Adverse Reactions:

Common:

- Dizziness
- Headache
- Dry cough
- Fatigue
- Increased urination
- Nausea
- Electrolyte disturbances

Less Common but Serious:

- **Angioedema**
- Severe hypotension
- Hyperkalemia or significant electrolyte imbalance
- Acute kidney injury
- Severe allergic reactions
- Pancreatitis
- Significant skin reactions

Seek immediate medical attention for swelling of the face, lips, tongue, or throat; difficulty breathing; fainting; or severe weakness.

Drug Interactions:

- **Potassium supplements and potassium-sparing diuretics** may increase the risk of hyperkalemia.
- **NSAIDs** may reduce the antihypertensive effect and increase the risk of renal impairment.
- **Lithium:** Hydrochlorothiazide and ramipril may increase lithium concentrations and toxicity risk.
- Other antihypertensive medicines may produce additive hypotension.
- **Antidiabetic medicines** may require monitoring because hydrochlorothiazide can affect blood glucose.
- **Aliskiren** may increase the risk of renal impairment, hyperkalemia, and hypotension in susceptible patients.
- **Sacubitril/valsartan:** An appropriate washout interval is required after an ACE inhibitor because of increased angioedema risk.
- Other medicines affecting electrolytes or renal function should be reviewed by the physician.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture and direct sunlight.
- Keep the tablets in their original packaging.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. **Rampolite H Tablet should be used under the supervision of a registered medical practitioner.** Blood pressure, renal function, and serum electrolytes should be monitored periodically during therapy.

Important: Ramipril is **contraindicated during pregnancy** because ACE inhibitors can cause serious fetal toxicity. Patients who become pregnant while taking this medicine should contact their physician promptly.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for