Cadray™ Lifecare

Repacept 2

Repaglinide 2mg

Effective Postprandial Control
Repaglinide effectively lowers post-meal blood glucose levels.

Rapid Onset of Action
Starts working within 30 minutes of administration.

Improves Glycemic Control
Helps achieve and maintain target HbA1c levels effectively.

Low Risk of Hypoglycemia
Lower risk when used appropriately with meals.

Flexible Dosing
Can be dosed with meals to suit individual patient needs.

Well Tolerated & Safe
Favorable safety profile for long-term management of Type 2 Diabetes.



Indications

- Type 2 Diabetes Mellitus
- Postprandial Hyperglycemia
- Adjunct to Diet & Exercise
- Glycemic Control
- Long Term Management

References:

1. Nauck MA, Meier JJ. Repaglinide: a review of its use in the management of type 2 diabetes mellitus. Drugs. 2002;62(12):1749–1768.
2. Owens DR. Repaglinide: a review of its clinical efficacy in the management of type 2 diabetes mellitus. Clin Ther. 2001;23(12):1925–1943.
3. Scheen AJ. Clinical pharmacokinetics of repaglinide. Clin Pharmacokinet. 2000;39(6):451–480.

WHO

Good Manufacturing Practice Certification • GMP Quality