Respihub™



Rifamail IS Tablets

Rifampicin IP 450 mg + Isoniazid IP 300 mg

The Powerful Duo Against Tuberculosis.

Proven Anti-TB Combination
Rifampicin and Isoniazid combination is the cornerstone of first-line therapy for tuberculosis. [1]

Bactericidal Synergy
Dual action targets actively dividing and dormant Mycobacterium tuberculosis for enhanced efficacy. [2]

Prevents Drug Resistance
Combination therapy minimizes the risk of emergence of drug-resistant TB strains. [3]

Shorter Treatment Duration
Enables effective treatment with shorter regimens and improved patient adherence. [4]

Better Patient Outcomes
Improves sputum conversion rates and reduces relapse rates in TB patients. [5]

Indications

Pulmonary Tuberculosis (PTB) | Extrapulmonary Tuberculosis (EPTB) | Drug Sensitive Tuberculosis | Sputum Positive TB Cases

Referenons:

1. World Health Organization. WHO treatment guidelines for drug-resistant Tuberculosis. Update 2022.
2. Grosset J. The sterilizing effect of rifampicin in experimental short-course chemotherapy of tuberculous infection. Tubercle. 1980;61(4):219-228.
3. Mitchison DA. Basic mechanisms of chemotherapy. Chest. 1979;76(6):771-781.
4. Nahid P, et al. Official American Thoracic Society/Centers for Disease Control and Prevention/Infectious Diseases Society of America Clinical Practice Guidelines: Treatment of Drug-Susceptible Tuberculosis. Clin Infect Dis. 2016;63(7):e147-e195.
5. Vynnycky E, Fine PEM. The natural history of tuberculosis: the implications of age-dependent risks of disease and the role of reinfection. Epidemiol Infect. 2000;124(2):183-201.