Information For the Use of
Rivaroxaban (20mg) Tablet

Rivabridge 20 Tablet

Rivabridge 20 Tablets

Composition:

Each tablet contains:

- **Rivaroxaban IP** – 20 mg

Pharmacological Action:

Rivaroxaban is a **direct oral anticoagulant (DOAC)** that selectively inhibits Factor Xa, an important enzyme in the blood coagulation pathway.

- **Factor Xa Inhibition:** Directly inhibits free and clot-bound Factor Xa.
- **Reduced Thrombin Generation:** Decreases thrombin formation and subsequent fibrin development.
- **Anticoagulant Action:** Helps prevent the formation and propagation of blood clots.
- **Thromboembolic Protection:** Helps reduce the risk of stroke and systemic embolism and treat venous thromboembolic disorders in appropriately selected patients.

Indications:

Rivabridge 20 Tablets is indicated for the management of:

- **Treatment of Deep Vein Thrombosis (DVT)**
- **Treatment of Pulmonary Embolism (PE)**
- **Reduction in the Risk of Recurrent DVT and PE**
- **Prevention of Stroke and Systemic Embolism in Non-Valvular Atrial Fibrillation**
- Other approved thromboembolic indications as determined by the treating physician.

Dosage and Administration:

- **Recommended Dose:** Dosage and duration depend on the indication, renal function, and individual patient factors and should be determined by the physician.
- **Administration:** The **20 mg tablet should be taken with food** to ensure adequate absorption.
- Swallow the tablet whole with water.
- Take the medicine at approximately the same time each day.
- Do not discontinue treatment abruptly without medical advice, as this may increase the risk of thrombotic events.

Precautions:

- Assess **renal and hepatic function** before and during treatment when clinically appropriate.
- Rivaroxaban increases the risk of bleeding. Report unusual bruising, prolonged bleeding, blood in urine or stools, or severe headache immediately.
- Use cautiously in patients with conditions associated with an increased risk of bleeding.
- Inform the physician before surgery, dental procedures, or invasive procedures.
- Avoid unnecessary use of **NSAIDs, antiplatelet agents, or other anticoagulants** unless prescribed.
- **Pregnancy and Lactation:** Use only when specifically advised by a physician.

Contraindications:

- Active clinically significant bleeding.
- Known hypersensitivity to rivaroxaban or any component of the formulation.
- Conditions associated with a high risk of major bleeding.
- Severe hepatic disease associated with coagulopathy and clinically relevant bleeding risk.
- Concomitant use with certain strong **CYP3A4 and P-glycoprotein inhibitors or inducers**, where clinically contraindicated.

Adverse Reactions:

Common:

- Bleeding
- Easy bruising
- Nosebleed
- Anemia
- Gastrointestinal discomfort

Less Common but Serious:

- Major bleeding
- Gastrointestinal bleeding
- Hematuria
- Intracranial bleeding
- Severe hypersensitivity reactions

Patients should seek immediate medical attention for uncontrolled bleeding, severe headache, weakness, dizziness, or blood in vomit, urine, or stools.

Drug Interactions:

- Concomitant use with **anticoagulants, antiplatelet agents, or NSAIDs** may increase bleeding risk.
- Strong **CYP3A4 and P-glycoprotein inhibitors** may increase rivaroxaban exposure and bleeding risk.
- Strong CYP3A4/P-glycoprotein inducers may decrease rivaroxaban exposure and reduce its anticoagulant effect.
- Inform the physician about all prescription medicines, OTC medicines, and herbal supplements being used.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture and direct sunlight.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. Rivabridge 20 Tablets should be used under the supervision of a registered medical practitioner. The dosage and duration should be individualized according to the patient's indication, renal function, bleeding risk, and overall clinical condition.

Manufactured in India for:

Cafoli™ LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for