Information For the Use of
Roxithromycin (150mg) Tablets

Roxyfoli 150 Tablet

ROXYFOLI 150 TABLET

Roxithromycin (150 mg)

1. Composition

Each film-coated tablet contains:
Roxithromycin 150 mg
Excipients q.s.

2. Pharmacological Action

Roxithromycin is a **macrolide antibiotic**. It inhibits bacterial protein synthesis by binding to the 50S subunit of bacterial ribosomes, thereby suppressing the growth of susceptible bacteria.

3. Indications

Roxithromycin is indicated for the treatment of susceptible bacterial infections, including:

- Upper respiratory tract infections such as **pharyngitis, tonsillitis and sinusitis**.
- Lower respiratory tract infections such as **acute bronchitis and community-acquired pneumonia**.
- Skin and soft-tissue infections.
- Certain genitourinary infections caused by susceptible organisms.

Use should be based on the suspected or confirmed bacterial pathogen and local antimicrobial-susceptibility patterns.

4. Dosage and Administration

Adults: The usual dose is **150 mg twice daily**, approximately every 12 hours, or as directed by the physician.

The tablet should generally be taken **at least 15 minutes before meals**, with sufficient water, and swallowed whole.

The duration of treatment depends on the type and severity of infection and should be determined by the physician.

5. Precautions

- Use only for infections caused by susceptible bacteria; antibiotics should not be used for viral infections such as uncomplicated common cold.
- Use cautiously in patients with **hepatic impairment**.
- Patients with a history of QT prolongation or significant cardiac rhythm disorders should be assessed before treatment.
- Use cautiously with medicines that may prolong the QT interval.
- Discontinue and seek medical attention if severe or persistent diarrhoea occurs, as antibiotic-associated colitis may develop.
- Complete the prescribed course unless otherwise directed by the physician.
- Monitor patients who develop signs of hypersensitivity or severe skin reactions.

6. Contraindications

Roxithromycin is contraindicated in patients with:

- Known hypersensitivity to **roxithromycin, erythromycin, other macrolides**, or any component of the formulation.
- Concomitant use with certain medicines where serious interactions are expected, particularly some **ergot alkaloids**.
- Conditions where use is contraindicated according to the approved product information.

7. Adverse Reactions

Possible adverse reactions include:

- Nausea, vomiting and abdominal discomfort
- Diarrhoea
- Headache and dizziness
- Abnormal taste
- Skin rash or itching
- Elevated liver enzymes

Rare but serious reactions may include **severe hypersensitivity, hepatotoxicity, QT prolongation and serious antibiotic-associated diarrhoea**.

8. Drug Interactions

Roxithromycin may interact with certain medicines, including:

- Drugs associated with **QT prolongation**, potentially increasing the risk of cardiac arrhythmias.
- Certain ergot alkaloids.
- Warfarin and other anticoagulants may require appropriate monitoring.
- Digoxin and other medicines whose concentrations may be affected by macrolide therapy.

The physician should be informed about all prescription, OTC and herbal medicines being taken.

9. Storage

Store below **30°C**, protected from light and moisture.

Keep out of reach of children.

10. Note / Important

Roxithromycin is an **antibacterial medicine** and should be used only when a bacterial infection is suspected or confirmed to be susceptible.

Do not share antibiotics with others or use leftover antibiotics. Complete the prescribed course as directed by the physician.

For use under medical supervision.

Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for