Critsila™



Silocaine
Injection

Lidocaine Hydrochloride Injection IP 1%

Fast Onset. Reliable Anesthesia. Maximum Safety.

Rapid Onset of Action
Provides quick and effective local anesthesia for procedural success.[1]

Effective Local Anesthesia
Ideal for infiltration, nerve block and topical anesthesia.[2]

Well Tolerated
Low incidence of adverse effects when used as directed.[3]

Versatile Use
Suitable for a wide range of surgical, diagnostic and therapeutic procedures.[4]

Proven Safety Profile
Extensively used with a long-standing record of safety and efficacy.[5]

Indications

Local Infiltration Anesthesia | Peripheral Nerve Blocks | Regional Anesthesia | Topical Anesthesia | Minor Surgical Procedures

References:

1. Covino BG, Vassalo HG. Local anesthetics: mechanisms of action and clinical use. N Engl J Med. 1976;294(24):1265-1269.
2. Neal JM, Barrington MJ, Brull R, et al. The second ASRA practice advisory on neurologic complications in regional anesthesia and pain medicine. Reg Anesth Pain Med. 2015;40(5):401-430.
3. Becker DE. Local anesthetics: review of pharmacological considerations. Anesth Prog. 2010;57(2):90-101.
4. Brown DL. Local and regional anesthesia. In: Butterworth JF, Mackey DC, Wasnick JD, eds. Morgan & Mikhail's Clinical Anesthesiology. 6th ed. McGraw-Hill; 2018:280-312.
5. Kopacz DJ, Neal JM. Current concepts in local anesthetic pharmacology. Anesthesiology. 2001;95(1):217-230.

For healthcare professionals only. Refer full prescribing information before use.