Information For the Use of
Sucralfate (1gm) Tablet

Sucraziff Tablet

Sucraziff Tablet

Composition:

Each tablet contains:

- **Sucralfate IP** – 1 g

Pharmacological Action:

Sucralfate is a **gastroprotective (anti-ulcer) agent** that acts locally on the gastric and duodenal mucosa. In the acidic environment of the stomach, it forms a viscous, adhesive protective barrier over ulcerated and inflamed tissues, shielding them from the damaging effects of gastric acid, pepsin, and bile salts. It promotes mucosal healing while having minimal systemic absorption.

- **Mucosal Protection:** Forms a protective coating over gastric and duodenal ulcers.
- **Ulcer Healing:** Promotes healing by protecting damaged mucosa from acid and pepsin.
- **Cytoprotective Effect:** Stimulates local production of prostaglandins and bicarbonate, enhancing mucosal defense.
- **Minimal Systemic Absorption:** Acts locally within the gastrointestinal tract with limited systemic exposure.

Indications:

Sucraziff Tablet is indicated for the treatment and prevention of acid-related gastrointestinal disorders, including:

- **Duodenal Ulcer**
- **Gastric Ulcer**
- **Gastritis**
- **Gastroesophageal Reflux Disease (GERD)**
- **Stress Ulcer Prophylaxis** (as advised by the physician)

Dosage and Administration:

- **Recommended Dose:** As directed by the physician.
- **Administration:** Take the tablet on an **empty stomach**, preferably **1 hour before meals and at bedtime**, with sufficient water.

- Avoid taking antacids within **30 minutes** before or after Sucralfate unless advised by the physician.
- Complete the prescribed course even if symptoms improve.

Precautions:

- Use with caution in patients with **renal impairment**, particularly during prolonged therapy.
- Maintain an interval of at least **2 hours** between Sucralfate and other oral medications, as it may reduce their absorption.
- Pregnancy and lactation: Use only under medical supervision if clearly indicated.
- Long-term therapy should be used only as recommended by the physician.

Contraindications:

- Known hypersensitivity to **Sucralfate** or any component of the formulation.
- Use with caution in patients with severe renal dysfunction due to the risk of aluminum accumulation.

Adverse Reactions:

Common:

- Constipation
- Dry mouth
- Nausea
- Indigestion
- Flatulence

Less Common:

- Dizziness
- Skin rash
- Pruritus
- Hypersensitivity reactions (rare)

Patients should consult their healthcare provider if severe allergic reactions or persistent gastrointestinal symptoms occur.

Drug Interactions:

- Sucralfate may reduce the absorption of medications such as **fluoroquinolone antibiotics**, **tetracyclines**, **levothyroxine**, **digoxin**, **phenytoin**, and **warfarin**.
- Maintain a minimum **2-hour interval** between Sucralfate and other oral medications.
- Antacids should not be taken within **30 minutes** of Sucralfate administration unless advised by the physician.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture and direct sunlight.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. Sucraziff Tablet should be used only under the supervision of a registered medical practitioner. The dosage and duration of therapy should be individualized based on the patient's clinical condition and the treating physician's judgment.

Manufactured in India for:

Cafoli™ LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for