Information For the Use of
Timolol (5mg) + Brimonidine (2mg) Eye Drops

Brimshell T Eye Drops

Brand Name: Brimshell T Eye Drops
Dosage Form: Eye Drops
Composition:
Each ml contains:
• Timolol Maleate IP equivalent to Timolol 5 mg
• Brimonidine Tartrate IP equivalent to Brimonidine 2 mg
• Sterile ophthalmic vehicle q.s.

Description:
Brimshell T Eye Drops is a sterile ophthalmic preparation containing Timolol, a non-selective beta-adrenergic blocker, and Brimonidine, an alpha-2 adrenergic receptor agonist. It is indicated for the reduction of elevated intraocular pressure in patients with glaucoma or ocular hypertension. It helps in lowering intraocular pressure by reducing aqueous humor production and enhancing aqueous outflow.

Indications:
Indicated for the treatment of:
• Open-angle glaucoma
• Ocular hypertension
• Elevated intraocular pressure requiring combination therapy
It helps in reducing intraocular pressure and minimizing the risk of optic nerve damage associated with glaucoma.

Dosage & Administration:
Instill one drop into the affected eye(s) twice daily or as directed by the physician.
Wash hands before administration.
Avoid touching the dropper tip to any surface to prevent contamination.
If using other ophthalmic medications, maintain an interval of at least 5–10 minutes between applications.

Pharmacological Action:

- **Timolol** reduces intraocular pressure by decreasing the production of aqueous humor in the ciliary body.
- **Brimonidine** is an alpha-2 adrenergic agonist that decreases aqueous humor production and increases uveoscleral outflow.

Together, these mechanisms provide effective reduction of elevated intraocular pressure.

Side Effects:
Common side effects include eye irritation, burning sensation, dry eyes, blurred vision, ocular redness, and headache.
Systemic effects such as fatigue, dizziness, bradycardia, or dry mouth may occur in some patients.

Precautions & Warnings:
• For ophthalmic use only
• Use with caution in patients with asthma, chronic obstructive pulmonary disease, or cardiac conduction disorders
• Remove contact lenses before instillation and reinsert after 15 minutes
• Monitor intraocular pressure regularly during treatment
• Discontinue use and seek medical advice if severe ocular irritation occurs

---

Drug Interactions:
May interact with systemic beta-blockers, antihypertensive agents, calcium channel blockers, and CNS depressants.
Caution is advised when used with monoamine oxidase inhibitors (MAOIs) or tricyclic antidepressants.

Storage:
Store below 25°C in a cool, dry place.
Protect from light.
Keep the bottle tightly closed after use.
Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for