Information For the Use of
Caroverine (20mg) Capsule

Otoverine 20 Capsule

Brand Name: Otoverine 20 Capsule
Dosage Form: Capsule
Composition:
Each capsule contains:
• Caroverine Hydrochloride IP equivalent to Caroverine 20 mg

Description:
Otoverine 20 Capsule contains Caroverine, an oto-neuroprotective agent with calcium channel-blocking and glutamate receptor antagonist properties. It is primarily indicated for the management of tinnitus (ringing in the ears) and certain inner ear disorders. Caroverine helps reduce abnormal excitatory neurotransmission within the auditory pathway, thereby alleviating tinnitus symptoms and improving auditory comfort. It may also provide relief in conditions associated with smooth muscle spasm.

Indications:
Indicated for the treatment of:
• Tinnitus (ringing or buzzing in the ears)
• Sensorineural tinnitus
• Vertigo associated with inner ear disorders (as advised by the physician)
• Adjunctive management of cochlear disorders
• Smooth muscle spasms in selected clinical conditions
It helps reduce the intensity of tinnitus, improve auditory comfort, and relieve smooth muscle spasm.

Dosage & Administration:
Dosage should be as directed by the physician.
The capsule should be swallowed whole with water and may be taken with or without food.
Continue therapy for the prescribed duration to obtain optimal clinical benefit.

Pharmacological Action:

- **Caroverine acts as an antagonist of excitatory glutamate receptors (including NMDA receptors), thereby reducing excessive neuronal excitation in the auditory pathway.**
- **It also exhibits calcium channel-blocking activity, contributing to smooth muscle relaxation and improved microcirculation in the inner ear.**
- **These actions help decrease abnormal auditory nerve activity associated with tinnitus and support inner ear function.**

Side Effects:
Common side effects include nausea, dizziness, headache, dry mouth, blurred vision, skin rash, itching, and mild gastrointestinal discomfort.
Rarely, hypersensitivity reactions or urinary retention may occur.

Precautions & Warnings:
• Use only under medical supervision.
• Use with caution in patients with hepatic impairment or gastrointestinal disorders.
• Avoid use in patients with known hypersensitivity to Caroverine.

• Safety during pregnancy and lactation has not been established; use only if clearly indicated by the physician.
• If dizziness occurs, avoid driving or operating machinery.

Drug Interactions:
No clinically significant drug interactions have been well established.
Use cautiously with other centrally acting medications or drugs affecting the nervous system. Inform the physician about all concomitant medications before initiating therapy.

Storage:
Store below 25°C in a cool, dry place.
Protect from light and moisture.
Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for