Information For the Use of
Tobramycin (0.3% w/w) Eye Ointment

Tobraxyle Eye Ointment

Brand Name: Tobraxyle Eye Ointment
Dosage Form: Eye Ointment
Composition:
Each gram contains:
• Tobramycin IP 0.3% w/w
• Ophthalmic ointment base q.s.

Description:
Tobraxyle Eye Ointment contains Tobramycin, an aminoglycoside antibiotic indicated for the treatment of external ocular infections caused by susceptible bacteria. It helps in eliminating pathogenic microorganisms, reducing infection-related symptoms, and promoting healing of affected ocular tissues. The ointment formulation provides prolonged contact with the ocular surface, enhancing local antibacterial activity.

Indications:
Indicated for the treatment of:
• Bacterial conjunctivitis
• Blepharitis
• Blepharoconjunctivitis
• Keratitis caused by susceptible organisms
• Other external ocular infections as determined by the physician
It helps in controlling bacterial growth, relieving ocular discomfort, and preventing the progression of infection.

Dosage & Administration:
Apply a small amount of ointment inside the lower conjunctival sac of the affected eye(s) as directed by the physician.
Wash hands thoroughly before application.
Avoid touching the tube tip to the eye or any surface to prevent contamination.
Complete the full course of treatment as prescribed.

Pharmacological Action:

- **Tobramycin binds to the 30S bacterial ribosomal subunit, interfering with protein synthesis and leading to bacterial cell death.**
- **It exhibits bactericidal activity against a broad range of susceptible Gram-positive and Gram-negative organisms.**

These actions help in effective eradication of bacterial eye infections.

Side Effects:

Common side effects include temporary blurred vision, eye irritation, burning sensation, redness, itching, and eyelid discomfort.
Rarely, hypersensitivity reactions or localized allergic responses may occur.

Precautions & Warnings:
- **For ophthalmic use only**
- **Not effective against viral or fungal eye infections**
- **Avoid contamination of the tube tip**
- **Prolonged use may result in overgrowth of non-susceptible organisms**
- **Discontinue use and seek medical advice if signs of hypersensitivity occur**
- **Contact lenses should generally not be worn during active ocular infection**

Drug Interactions:
No significant systemic drug interactions are expected due to minimal systemic absorption.
If used with other ophthalmic medications, maintain an appropriate interval between administrations.

Storage:
Store below 25°C in a cool, dry place.
Protect from light.
Do not freeze.
Keep the tube tightly closed after use.
Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for