Information For the Use of
Torasemide (40mg) Tablet

Torslide 40 Tablet

Brand Name: Torslide 40 Tablet
Dosage Form: Tablet
Composition:
Each tablet contains:
• Torsemide IP 40 mg

Description:
Torslide 40 Tablet contains Torsemide, a potent loop diuretic indicated for the management of edema associated with congestive heart failure, chronic kidney disease, hepatic cirrhosis, and hypertension. Torsemide inhibits sodium and chloride reabsorption in the thick ascending limb of the loop of Henle, promoting diuresis, reducing fluid overload, and lowering blood pressure. It provides effective and sustained diuretic action with predictable oral bioavailability.

Indications:
Indicated for the treatment of:
• Edema associated with congestive heart failure
• Edema due to chronic kidney disease
• Edema associated with hepatic cirrhosis
• Essential hypertension
• Fluid overload requiring loop diuretic therapy
It helps remove excess fluid from the body, reduces swelling, improves symptoms of fluid retention, and assists in controlling blood pressure.

Dosage & Administration:
Dosage should be individualized and administered as directed by the physician.
The tablet may be taken with or without food, preferably at the same time each day.
To minimize nocturnal urination, it is advisable to take the tablet in the morning.
Regular monitoring of renal function, electrolytes, blood pressure, and fluid balance is recommended during therapy.

Pharmacological Action:

- **Torsemide is a loop diuretic that inhibits the sodium-potassium-chloride ($Na^+/K^+/2Cl^-$) cotransporter in the thick ascending limb of the loop of Henle.**
- **This inhibition increases the excretion of sodium, chloride, water, potassium, calcium, and magnesium, resulting in effective diuresis.**
- **By reducing plasma volume and extracellular fluid, Torsemide decreases edema, lowers blood pressure, and reduces cardiac preload in patients with heart failure.**

Side Effects:
Common side effects include increased urination, dizziness, headache, dehydration, muscle cramps, weakness, hypotension, nausea, and electrolyte disturbances such as hypokalemia and hyponatremia. Rarely, hyperuricemia, hearing disturbances, renal impairment, hypersensitivity reactions, or severe electrolyte imbalance may occur.

Precautions & Warnings:

• Monitor serum electrolytes, renal function, and blood pressure regularly during treatment.
• Use with caution in patients with renal impairment, hepatic dysfunction, diabetes mellitus, gout, or electrolyte abnormalities.
• Correct dehydration and electrolyte imbalance before initiating therapy whenever possible.
• Patients should avoid sudden changes in posture to minimize orthostatic hypotension.
• Use during pregnancy and lactation only if clearly indicated by the physician.

Drug Interactions:
May interact with antihypertensive agents, aminoglycoside antibiotics, NSAIDs, lithium, digoxin, corticosteroids, antidiabetic medications, and other diuretics. Concomitant use may increase the risk of hypotension, nephrotoxicity, ototoxicity, or electrolyte imbalance; appropriate monitoring is recommended.

Storage:
Store below 25°C in a cool, dry place.
Protect from light and moisture.
Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for