Information For the Use of
Timolol (5mg) + Travoprost (40mcg) Eye Drop

Travopine T Eye Drop

Travopine T Eye Drops

Composition:

Each mL contains:

- **Timolol** – 5 mg
- **Travoprost** – 40 mcg

Pharmacological Action:

Travopine T combines a **beta-adrenergic blocker** with a **prostaglandin analogue** to reduce elevated intraocular pressure (IOP).

- **Timolol:** Blocks β_1 and β_2 adrenergic receptors in the ciliary body, reducing aqueous humour production.
- **Travoprost:** Increases uveoscleral outflow of aqueous humour, thereby lowering IOP.
- **Combined Action:** The complementary mechanisms provide greater IOP reduction than either component alone in appropriately selected patients.
- **Pressure Control:** Helps maintain IOP within the target range and reduces the risk of progressive optic nerve damage associated with glaucoma.

Indications:

Travopine T Eye Drops is indicated for the reduction of elevated intraocular pressure in:

- **Primary Open-Angle Glaucoma**
- **Ocular Hypertension**
- Patients requiring additional IOP reduction when monotherapy provides inadequate control

---

Dosage and Administration:

- **Recommended Dose:** Instill **1 drop into the affected eye(s) once daily**, or as directed by the ophthalmologist.
- Preferably administer at approximately the **same time each day**.
- Do not exceed **one dose per day**, as more frequent administration of travoprost may reduce the IOP-lowering effect.
- Wash hands before use.
- Tilt the head back and gently pull down the lower eyelid.

- Instill one drop without touching the dropper tip to the eye or surrounding surfaces.
- Close the eye gently and apply **nasolacrimal occlusion for approximately 1–2 minutes** to reduce systemic absorption.
- If more than one ophthalmic medicine is prescribed, allow an appropriate interval between applications.
- Remove contact lenses before administration and generally wait **at least 15 minutes** before reinserting them.

Precautions:

- For **ophthalmic use only**.
- Timolol may undergo systemic absorption and can cause **bradycardia, hypotension, bronchospasm, or conduction abnormalities** in susceptible patients.
- Use cautiously in patients with **cardiac disease, diabetes, thyroid disorders, or peripheral vascular disease**.
- Travoprost may cause **increased iris pigmentation**, eyelash growth, and changes in eyelid/periocular pigmentation. Iris pigmentation may be permanent.
- Prostaglandin analogues may rarely cause **macular edema**, particularly in patients with aphakia, pseudophakia with a torn posterior lens capsule, or other risk factors.
- Use cautiously in patients with a history of **uveitis or ocular inflammation**.
- Do not allow the bottle tip to contact the eye, eyelids, or other surfaces.
- **Pregnancy and Lactation:** Use only when clearly indicated and under specialist supervision.

Contraindications:

- Known hypersensitivity to **timolol, travoprost, or any component of the formulation**.
- **Bronchial asthma or a history of bronchial asthma**.
- Severe chronic obstructive pulmonary disease with significant bronchospastic disease.
- **Sinus bradycardia**.
- Second- or third-degree **atrioventricular block** without a functioning pacemaker.
- **Overt cardiac failure or cardiogenic shock**.

Adverse Reactions:

Common:

- Eye redness
- Eye irritation or discomfort
- Burning or stinging
- Dry eye
- Blurred vision
- Increased eyelash growth
- Changes in iris or eyelid pigmentation

Systemic Effects:

- Bradycardia
- Hypotension
- Fatigue
- Dizziness
- Shortness of breath

Rare but Serious:

- Bronchospasm
- Cardiac conduction disturbances
- Severe bradycardia
- Macular edema
- Severe hypersensitivity reactions

Seek immediate medical attention for difficulty breathing, fainting, severe dizziness, marked reduction in heart rate, or sudden visual changes.

Drug Interactions:

- Other **topical or systemic beta-blockers** may produce additive effects on heart rate and blood pressure.
- **Calcium-channel blockers**, cardiac glycosides, and other medicines affecting cardiac conduction may increase the risk of bradycardia or heart block.
- **Catecholamine-depleting medicines** may enhance hypotension or bradycardia.
- **Systemic CYP2D6 inhibitors** may increase systemic beta-blocking effects of timolol.
- Other glaucoma medicines may have additive IOP-lowering effects.
- If multiple eye drops are prescribed, administer them separately with an appropriate interval.

Storage:

- Store in a cool, dry place below **25°C**, or according to the manufacturer's approved storage conditions.
- Protect from direct sunlight and excessive heat.
- Keep the bottle tightly closed when not in use.
- Do not touch the dropper tip to any surface.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. **Travopine T Eye Drops should be used under the supervision of a qualified ophthalmologist**, with periodic monitoring of intraocular pressure.

Important: Because timolol can be systemically absorbed after topical ocular administration, patients with a history of **asthma, COPD, bradycardia, heart block, or cardiac failure** should be carefully evaluated before treatment.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for