Information For the Use of
Ursodeoxycholic Acid (300mg) Tablets

Udostep 300 Tablet

UDOSTEP 300 TABLETS

Composition

Each film-coated tablet contains:

Ursodeoxycholic Acid I.P. 300 mg
Excipients … q.s.

Pharmacological Action

Ursodeoxycholic Acid (UDCA) is a hydrophilic bile acid that modifies the composition and flow of bile. It decreases the relative proportion of hydrophobic, potentially hepatotoxic bile acids and promotes bile secretion.

UDCA exerts **hepatoprotective and cytoprotective effects** on hepatocytes and cholangiocytes. It also reduces biliary cholesterol saturation and may facilitate the gradual dissolution of appropriately selected cholesterol-rich gallstones.

Indications

Udostep 300 Tablets may be indicated for:

- Primary biliary cholangitis (PBC)
- Cholestatic liver diseases where UDCA therapy is appropriate
- Selected cholesterol-rich, radiolucent gallstones for medical dissolution
- Other approved hepatobiliary conditions as directed by the physician

The exact indication should follow the approved product-specific prescribing information.

Dosage and Administration

Adults:
Dosage is individualized according to the indication, body weight and clinical response.

For **primary biliary cholangitis**, a commonly used dosage is approximately **13–15 mg/kg/day**, administered in divided doses according to the physician's prescription.

For gallstone dissolution, dosage is generally calculated according to **body weight** and administered as prescribed.

Swallow the tablet whole with water. When prescribed in divided doses, administration with food may improve tolerability.

Treatment duration depends on the indication and may extend for several months or longer.

Monitoring:
Periodic monitoring of liver-function parameters, including **ALT, AST, alkaline phosphatase, γ-GT and bilirubin**, is recommended during treatment.

Precautions

- Use under medical supervision in patients with hepatic or biliary disease.
- Perform appropriate liver-function monitoring during treatment.
- Treatment should be reassessed if liver biochemical parameters deteriorate significantly.
- Before treatment for gallstone dissolution, appropriate imaging should confirm that the stones are suitable for medical dissolution.
- Not all gallstones respond to UDCA therapy.
- Use cautiously in patients with impaired bile flow or biliary obstruction.
- Persistent abdominal pain, biliary colic, jaundice or worsening liver-function abnormalities require medical evaluation.

Contraindications

Udostep 300 Tablets are contraindicated in patients with:

- Hypersensitivity to ursodeoxycholic acid or any component of the formulation.
- Complete biliary obstruction.
- Acute inflammation of the gallbladder or biliary tract when bile drainage is impaired.
- Non-functioning gallbladder when UDCA is being used specifically for gallstone dissolution.
- Calcified or otherwise unsuitable gallstones for medical dissolution.

Adverse Reactions

Common/possible:

- Diarrhoea or loose stools
- Abdominal discomfort
- Abdominal pain
- Nausea
- Dyspepsia

Other possible reactions:

- Vomiting
- Pruritus
- Skin rash

- Rare worsening of liver-function abnormalities

Persistent or severe diarrhoea should be reported to the treating physician.

Drug Interactions

Cholestyramine and Colestipol:
May reduce absorption of ursodeoxycholic acid. If concomitant therapy is required, the medicines should be administered at appropriately separated times.

Aluminium-containing antacids:
May reduce the absorption of UDCA and should not be taken simultaneously.

Other medicines affecting bile-acid metabolism:
Concomitant medicines should be reviewed by the physician to assess potential effects on UDCA absorption or efficacy.

Patients should inform their healthcare professional about all prescription, OTC and herbal medicines being used.

Pregnancy and Lactation

Use during pregnancy should be based on a medical assessment of the potential benefits and risks. UDCA is used therapeutically in selected cholestatic liver disorders during pregnancy under specialist supervision.

Use during breastfeeding should likewise be based on medical advice.

Storage

Store in a **cool, dry place**, protected from excessive heat, moisture and direct sunlight.

Keep the tablets in their original packaging and **out of reach of children**.

Note

For use under medical supervision.

Udostep 300 Tablets should be used according to the indication, body weight and clinical response. Regular clinical assessment and liver-function monitoring are recommended during prolonged treatment.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for