Information For the Use of
Valacyclovir (500mg) Tablets

Valaclo 500 Tablets

VALACLO 500 TABLET

Valacyclovir IP (500 mg)

1. Composition

Each film-coated tablet contains:
Valacyclovir Hydrochloride equivalent to Valacyclovir 500 mg
Excipients q.s.

2. Pharmacological Action

Valacyclovir is the **L-valyl ester prodrug of acyclovir**. Following oral administration, it is rapidly converted to acyclovir, which is phosphorylated by viral and cellular enzymes to its active triphosphate form.

Acyclovir triphosphate inhibits viral DNA polymerase and terminates viral DNA replication. It is active primarily against **Herpes simplex virus (HSV)** and **Varicella-zoster virus (VZV)**.

3. Indications

Valacyclovir is indicated for the treatment and management of susceptible herpesvirus infections, including:

- **Herpes zoster (shingles)**.
- **Herpes simplex infections**, including genital herpes.
- Treatment of recurrent genital herpes.
- Suppression of recurrent genital herpes in appropriate patients.
- Reduction of transmission of genital herpes in selected serodiscordant couples, when used as part of an appropriate management strategy.

4. Dosage and Administration

The dosage and duration depend on the indication, renal function and clinical condition.

Adults: Common regimens include **500 mg or 1 g at appropriate intervals**, depending on the indication and treatment phase.

Treatment should generally be started as early as possible after the onset of symptoms.

The tablet may be taken **with or without food** and should be swallowed with sufficient water.

Patients with renal impairment require dose adjustment.

5. Precautions

- Adequate hydration should be maintained, particularly in elderly patients and those at risk of renal impairment.
- Dose adjustment is required in patients with **renal impairment**.
- Use cautiously in elderly patients because of increased risk of renal and neurological adverse effects.
- Patients receiving high doses or prolonged treatment should be appropriately monitored.
- Rare neurological reactions such as confusion, hallucinations, agitation or seizures may occur, particularly in patients with renal impairment receiving excessive doses.
- Valacyclovir does not cure herpes infection and does not eliminate the risk of transmission.
- Patients with genital herpes should follow appropriate safer-sex practices and medical advice regarding transmission prevention.

6. Contraindications

Valacyclovir is contraindicated in patients with:

- Known hypersensitivity to **valacyclovir, acyclovir, or any component of the formulation**.
- Previous clinically significant hypersensitivity reaction to valacyclovir or acyclovir.

7. Adverse Reactions

Common adverse reactions may include:

- Headache
- Nausea
- Abdominal discomfort
- Vomiting
- Diarrhoea

Less common or serious reactions may include:

- Renal impairment or acute kidney injury
- Neurological effects such as confusion, hallucinations or seizures
- Severe hypersensitivity reactions
- Serious skin reactions, rarely

8. Drug Interactions

- Concomitant use with other **nephrotoxic medicines** may increase the risk of renal impairment.
- Caution is advised with medicines that may affect renal function or renal tubular secretion.
- Patients receiving other medicines that can cause neurological adverse effects should be monitored appropriately.

Inform the physician about all prescription, OTC and herbal medicines being taken.

9. Storage

Store below **30°C**, protected from light and moisture.

Keep out of reach of children.

10. Note / Important

Valacyclovir is an **antiviral medicine** and is most effective when treatment is initiated promptly after symptoms begin.

It controls herpesvirus infections but does not eradicate latent virus from the body. The dose must be adjusted according to **renal function**, particularly in elderly or renally impaired patients.

For use under medical supervision.

Keep out of reach of children.

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for