Vasohype 20 Injection

VASOHYPE 20 INJECTION

Composition

Each vial/ampoule contains:

Vasopressin 20 IU
Excipients … q.s.

For intravenous use after appropriate dilution, as applicable to the product presentation.

Pharmacological Action

Vasopressin is a naturally occurring peptide hormone with potent vasoconstrictor and antidiuretic activity.

It acts primarily through **V1a receptors** on vascular smooth muscle, producing vasoconstriction and increasing systemic vascular resistance and arterial blood pressure.

Vasopressin also acts through **V2 receptors** in the renal collecting ducts, promoting water reabsorption.

In vasodilatory shock, vasopressin can increase vascular tone and blood pressure and may be used as an adjunct to catecholamine vasopressors.

Indications

Vasohype 20 Injection may be used, under specialist/critical-care supervision, for:

- **Vasodilatory shock with persistent hypotension** despite adequate fluid resuscitation and catecholamine therapy.
- Vasopressor support in selected cases of **septic shock**.
- Vasopressor support in selected cases of **post-cardiotomy vasodilatory shock**.

Vasopressin should be used as part of comprehensive haemodynamic management and not as a substitute for appropriate fluid resuscitation and treatment of the underlying cause of shock. FDA prescribing information specifically identifies vasodilatory shock in adults with persistent hypotension despite fluids and catecholamines as an indication.

Dosage and Administration

Adults:

Vasopressin dosing must be individualized according to the clinical condition, blood pressure and response to therapy.

For **septic shock**, commonly used infusion rates are in the range of:

0.01–0.07 units/minute IV, according to clinical response and the applicable prescribing protocol.

For **post-cardiotomy shock**, infusion rates may be in the range of:

0.03–0.1 units/minute IV, under specialist supervision.

Administration:

- Administer by **controlled intravenous infusion** using an appropriate infusion device.
- The 20 units/mL presentation should be appropriately diluted with a compatible diluent such as **0.9% Sodium Chloride Injection or 5% Dextrose Injection**, according to the approved product instructions.
- Continuous monitoring of blood pressure, cardiac rhythm, peripheral perfusion and urine output is recommended.
- Do not administer as an undiluted IV bolus unless specifically permitted by the approved product information for the particular presentation.

Precautions

- Vasopressin should be administered only by healthcare professionals experienced in **critical-care haemodynamic management**.
- Monitor arterial blood pressure and cardiac function continuously or at appropriate frequent intervals.
- Excessive vasoconstriction may cause **myocardial, digital, mesenteric or other tissue ischemia**.
- Use cautiously in patients with significant **cardiac dysfunction or ischemic heart disease** because cardiac output may deteriorate.
- Monitor serum sodium and fluid balance during prolonged therapy.
- Abrupt discontinuation after prolonged therapy may be associated with **reversible diabetes insipidus**, with polyuria, dilute urine and hypernatremia.
- Assess peripheral and visceral perfusion during treatment.
- Correct significant hypovolaemia and treat the underlying cause of shock as appropriate.
- Use with caution in patients with vascular disease or conditions where excessive vasoconstriction may be harmful.

Contraindications

Vasohype 20 Injection is contraindicated in patients with:

- Known hypersensitivity to **vasopressin** or any component of the formulation.
- Additional contraindications related to any formulation-specific preservative or excipient should be considered according to the product's approved prescribing information.

Adverse Reactions

Common/Possible:

- Decreased cardiac output
- Bradycardia
- Tachyarrhythmias
- Hyponatremia
- Headache
- Abdominal cramps or nausea

Vascular/Ischemic effects:

- Peripheral ischemia
- Digital ischemia
- Skin ischemia
- Coronary ischemia
- Mesenteric ischemia

Other serious effects:

- Cardiac dysfunction
- Reduced perfusion of vital organs
- Reversible diabetes insipidus following discontinuation

The FDA label identifies decreased cardiac output, bradycardia, tachyarrhythmias, hyponatremia and ischemic events among the most important adverse reactions.

Drug Interactions

Catecholamines:
The vasopressor effects of vasopressin and catecholamines may be **additive**. Blood pressure and tissue perfusion should therefore be monitored closely.

Indomethacin:
May prolong or increase the effects of vasopressin.

Ganglionic blocking agents:
May increase the pressor response to vasopressin.

Medicines associated with SIADH:
May increase the pressor response and contribute to water retention or hyponatremia.

Medicines causing diabetes insipidus:
May reduce the pressor/antidiuretic response to vasopressin.

Patients should inform the treating physician about all prescription medicines, OTC medicines and other treatments being received.

Pregnancy and Lactation

Vasopressin may produce **uterine contractions** and should therefore be used during pregnancy only when clearly indicated and under specialist medical supervision.

There are limited data regarding use during lactation. The decision to administer vasopressin to a breastfeeding woman should be based on the clinical necessity and medical assessment.

Storage

Store according to the **product-specific approved storage conditions**.

Protect from excessive heat and light and keep the container in its original packaging until use.

Keep out of reach of children.

Note

For intravenous use only. To be administered by or under the supervision of a qualified healthcare professional.

Vasohype 20 Injection should be used only in appropriately monitored clinical settings. The exact **dilution, infusion concentration, dose and duration** should follow the approved product-specific prescribing information and institutional critical-care protocol.

Important: For final printed IP literature, verify whether the product is labelled as **20 IU per vial/ampoule or 20 IU/mL**, as this distinction is critical for preparation and administration.

Remove unsupported FDA referencesResolve the composition presentation
You're on the free plan
ChatGPT gets less accurate and may forget details in long conversations. Upgrade to chat longer with better memory.
Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for