Information For the Use of
Fluticasone Furoate (200mcg) + Vilanterol (25mcg) Inhalation

Flutialfa VT 200 Inhalation

FLUTIALFA VT 200 INHALATION

Composition

Each delivered dose contains:

Vilanterol 25 mcg
Fluticasone Furoate 200 mcg
Excipients … q.s.

Pharmacological Action

Fluticasone Furoate is a potent inhaled corticosteroid with anti-inflammatory activity. It acts locally in the airways to reduce airway inflammation, mucosal oedema and inflammatory responses associated with asthma.

Vilanterol is a long-acting selective β_2-adrenergic receptor agonist (LABA). Activation of β_2-receptors in bronchial smooth muscle increases intracellular cyclic AMP and produces prolonged **bronchodilation**.

The combination provides complementary **anti-inflammatory and long-acting bronchodilator activity**, improving airway calibre and helping maintain control of respiratory symptoms.

Indications

Flutialfa VT 200 Inhalation is indicated for the **regular maintenance treatment of asthma** in patients for whom treatment with an inhaled corticosteroid and long-acting β_2-agonist combination is appropriate.

It may be considered in patients:

- Not adequately controlled with an inhaled corticosteroid and an as-needed short-acting β_2-agonist
- Already adequately controlled with an inhaled corticosteroid and LABA
- Requiring maintenance therapy to improve asthma symptom control and reduce exacerbation risk

The 200/25 mcg strength is used as a higher-strength maintenance option in appropriate asthma patients.

Important: It is **not indicated for the relief of acute bronchospasm or sudden asthma attacks**. A suitable short-acting rescue bronchodilator should be available for acute symptoms.

Dosage and Administration

Adults:
The recommended dose of **Fluticasone Furoate 200 mcg + Vilanterol 25 mcg is one inhalation once daily**, administered by oral inhalation, when this strength is appropriate for the patient's asthma control.

- Administer at approximately the **same time each day**.
- Do not use more than **one inhalation in 24 hours**.
- Do not use an additional dose for acute breathlessness.
- If symptoms occur between doses, use the prescribed short-acting rescue inhaler.
- After inhalation, **rinse the mouth with water and spit it out** to reduce the risk of oral/oropharyngeal candidiasis.

The inhaler should be used with the correct inhalation technique according to the device-specific instructions.

Precautions

- **Not for acute relief:** Do not initiate or use this combination as rescue treatment during an acute severe asthma episode.
- LABA therapy should not be used as monotherapy for asthma; here, vilanterol is combined with an inhaled corticosteroid.
- Monitor patients for **oral candidiasis**; rinsing the mouth after inhalation is recommended.
- Use cautiously in patients with cardiovascular disorders, arrhythmias, hypertension or a history of QT prolongation.
- Use cautiously in patients with **hyperthyroidism, diabetes mellitus or seizure disorders**.
- β_2-agonists may produce **hypokalaemia** and transient increases in blood glucose in susceptible patients.
- Prolonged inhaled corticosteroid treatment may be associated with systemic corticosteroid effects, including effects on adrenal function, bone density, eyes and growth in susceptible populations.
- Patients with moderate or severe hepatic impairment may have increased systemic exposure to fluticasone furoate and require appropriate monitoring.

Contraindications

Flutialfa VT 200 Inhalation is contraindicated in:

- Hypersensitivity to **fluticasone furoate, vilanterol** or any component of the formulation.
- Severe hypersensitivity to **milk proteins**, where applicable to a lactose-containing dry-powder formulation.
- Primary treatment of **status asthmaticus** or other acute episodes of asthma requiring intensive measures.

Adverse Reactions

Common/possible:

- Headache
- Nasopharyngitis
- Upper respiratory tract infection
- Cough
- Hoarseness/dysphonia
- Oral or oropharyngeal candidiasis
- Tremor
- Palpitations
- Muscle cramps

Other/serious possible reactions:

- Paradoxical bronchospasm
- Tachycardia or cardiac arrhythmias
- Hypokalaemia
- Hyperglycaemia
- Hypersensitivity reactions
- Adrenal suppression with excessive systemic corticosteroid exposure
- Glaucoma or cataract with prolonged corticosteroid exposure

Drug Interactions

Strong CYP3A4 inhibitors:
Medicines such as ketoconazole, ritonavir and cobicistat may increase systemic exposure to fluticasone furoate and vilanterol. Use requires caution and appropriate monitoring.

β-blockers:
May antagonise the bronchodilator effect of vilanterol and may cause bronchospasm. Concomitant use should generally be avoided unless clinically necessary.

MAO inhibitors and tricyclic antidepressants:
May potentiate cardiovascular effects of β_2-agonists; use with caution.

Non-potassium-sparing diuretics:
May increase the risk of β_2-agonist-associated hypokalaemia and ECG changes.

Other medicines that prolong the QT interval should be used cautiously with vilanterol.

Pregnancy and Lactation

Use during pregnancy should be considered only when the expected benefit justifies the potential risk. A healthcare professional should assess the need for continued maintenance therapy.

Use during breastfeeding should similarly be based on an assessment of maternal benefit and potential risk to the infant.

Storage

Store the inhaler according to the **device-specific approved storage conditions**.

Keep in the original packaging, protected from moisture, heat and direct sunlight.

Keep out of reach of children.

Do not use after the expiry date or if the inhaler has been damaged.

Note

For use under medical supervision.

Flutialfa VT 200 Inhalation is a **maintenance controller medication and not a rescue inhaler**. It should be administered regularly as prescribed and should not be used more frequently than the recommended once-daily dose. Correct inhaler technique and regular follow-up are important for optimal asthma control.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for