Information For the Use of
Prednisolone (10mg) Eye Drop

Predslide Eye Drop

Predslide Eye Drops

Composition:

Each mL contains:

- **Prednisolone** – 10 mg

Important: Please verify the exact **prednisolone salt and concentration** from the approved product specification before finalizing the IP literature, as ophthalmic prednisolone formulations can differ in salt form and strength.

Pharmacological Action:

Prednisolone is a **corticosteroid** that suppresses inflammation in ocular tissues.

- **Anti-inflammatory Action:** Reduces the production and release of inflammatory mediators.
- **Reduction of Ocular Inflammation:** Helps decrease redness, swelling, irritation, and inflammatory cell infiltration.
- **Immunosuppressive Action:** Suppresses local immune responses associated with ocular inflammation.
- **Symptomatic Relief:** Helps relieve discomfort associated with steroid-responsive inflammatory eye conditions.

Indications:

Predslide Eye Drops is indicated for the treatment of **steroid-responsive ocular inflammation**, including:

- **Anterior Uveitis**
- **Iritis and Iridocyclitis**
- **Post-operative Ocular Inflammation**
- **Allergic and Inflammatory Eye Conditions**
- Other steroid-responsive inflammatory conditions of the eye, as prescribed by an ophthalmologist

---

Dosage and Administration:

- **Recommended Dose:** Instill the prescribed number of drops into the affected eye(s) as directed by the ophthalmologist.
- Wash hands before use.
- Tilt the head back and gently pull down the lower eyelid.
- Avoid touching the dropper tip to the eye, eyelids, or surrounding surfaces.
- Close the eye gently after instillation.
- If using other eye drops, maintain an appropriate interval between different preparations.
- Do not discontinue prolonged treatment abruptly without medical advice.

Precautions:

- Use under **ophthalmic supervision**, particularly for prolonged treatment.
- Prolonged corticosteroid use may increase **intraocular pressure** and the risk of glaucoma.
- Long-term use may contribute to **cataract formation**.
- Corticosteroids may delay corneal wound healing and can mask or worsen certain ocular infections.
- Use cautiously in patients with **glaucoma, ocular herpes infection, or a history of ocular infections**.
- **Pregnancy and Lactation:** Use only when clearly indicated and under medical supervision.

Contraindications:

- Known hypersensitivity to prednisolone or any component of the formulation.
- **Untreated acute purulent ocular infections**.
- Active fungal infections of the eye.
- **Herpes simplex keratitis** and other viral ocular infections where corticosteroid therapy is contraindicated.
- Mycobacterial infections of the eye unless specifically managed by an ophthalmologist.

Adverse Reactions:

Common:

- Temporary burning or stinging
- Mild eye irritation
- Temporary blurred vision after instillation

Less Common but Serious:

- Increased intraocular pressure

- Glaucoma
- Cataract formation with prolonged use
- Delayed corneal wound healing
- Secondary ocular infection
- Corneal or scleral thinning in susceptible patients
- Severe hypersensitivity reactions

Drug Interactions:

- Concurrent use with other **ophthalmic corticosteroids** may increase the risk of steroid-related adverse effects.
- If multiple ophthalmic preparations are prescribed, administer them separately with an appropriate interval.
- Inform the ophthalmologist about all prescription, OTC, and ophthalmic medicines being used.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from direct sunlight and excessive heat.
- Keep the container tightly closed when not in use.
- Avoid touching the dropper tip to any surface.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. Predslide Eye Drops should be used only under the supervision of a qualified ophthalmologist. Prolonged or inappropriate use of topical ocular corticosteroids may increase intraocular pressure and may mask or worsen ocular infections.

Because the product is labeled as prednisolone 10 mg per mL, the final literature should state the formulation as provisional until the approved salt and strength are verified to avoid an inaccurate composition claim.

Manufactured in India for:

Cafoli™ LIFECARE

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113

Corporate Office :

Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113

TM: Trademark Applied for