Information For the Use of
Prednisolone (5mg/5ml) Solution

Predplit Solution

Brand Name: Predplit Solution
Dosage Form: Oral Suspension
Composition:
Each 5 ml contains:
• Prednisolone IP 5 mg
• Flavoured suspension base q.s.

Description:
Predplit Solution contains Prednisolone, a synthetic glucocorticoid with potent anti-inflammatory, antiallergic, and immunosuppressive properties. It is indicated for the treatment of a wide range of inflammatory, allergic, autoimmune, and endocrine disorders. Prednisolone helps suppress inflammatory responses, reduce tissue swelling, and modulate immune activity, thereby providing symptomatic relief in acute and chronic conditions.

Indications:
Indicated for the treatment of:
• Allergic disorders (including allergic rhinitis and urticaria)
• Bronchial asthma and acute exacerbations
• Autoimmune disorders
• Rheumatic and musculoskeletal disorders
• Dermatological inflammatory conditions
• Nephrotic syndrome
• Inflammatory bowel diseases
• Endocrine disorders requiring corticosteroid therapy
It helps in reducing inflammation, controlling allergic reactions, and suppressing excessive immune responses.

Dosage & Administration:
Dosage should be individualized and administered strictly as directed by the physician.
Shake the bottle well before use.
Measure the prescribed dose using a calibrated measuring device.
The suspension is preferably taken after food to reduce gastric irritation.
Do not discontinue therapy abruptly, especially after prolonged treatment.

Pharmacological Action:

- **Prednisolone is a glucocorticoid that binds to intracellular corticosteroid receptors and regulates gene expression.**
- **It inhibits the production of inflammatory mediators such as prostaglandins, leukotrienes, and cytokines while suppressing leukocyte migration and capillary permeability.**
- **It also exerts immunosuppressive effects by reducing immune cell activity and inflammatory responses.**

These actions provide effective control of inflammatory, allergic, and autoimmune conditions.

Side Effects:
Common side effects include increased appetite, weight gain, indigestion, mood changes, insomnia, headache, and elevated blood glucose levels.
Prolonged therapy may lead to hypertension, osteoporosis, adrenal suppression, increased susceptibility to infections, cataracts, glaucoma, or Cushingoid features.

Precautions & Warnings:
• Use only under medical supervision.
• Do not discontinue treatment abruptly after prolonged use.
• Use with caution in patients with diabetes mellitus, hypertension, osteoporosis, peptic ulcer disease, glaucoma, hepatic impairment, or renal impairment.
• Monitor growth in pediatric patients receiving long-term therapy.
• Patients receiving prolonged corticosteroid therapy should be monitored for adrenal suppression and infections.

Drug Interactions:
May interact with NSAIDs, anticoagulants, antidiabetic agents, vaccines, rifampicin, phenytoin, phenobarbital, ketoconazole, cyclosporine, and diuretics.
Concurrent use with live vaccines should be avoided in immunosuppressed patients.

Storage:
Store below 25°C in a cool, dry place.
Protect from light.
Do not freeze.
Shake well before use.
Keep out of reach of children.

Manufactured in India for:

Cafoli™
LIFECARE

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for