Information For the Use of
Warfarin (1mg) Tablet

Warwise 1 Tablet

Warwise 1 Tablet

Composition:

Each tablet contains:

- **Warfarin** – 1 mg

Pharmacological Action:

Warfarin is an **oral anticoagulant** that reduces the formation of vitamin K-dependent clotting factors.

- **Vitamin K Antagonism:** Inhibits vitamin K epoxide reductase, reducing the regeneration of active vitamin K.
- **Reduced Clotting Factors:** Decreases synthesis of Factors II, VII, IX, and X, as well as anticoagulant proteins C and S.
- **Anticoagulant Action:** Prolongs blood clotting time and helps prevent formation and growth of harmful blood clots.
- **Thromboembolic Protection:** Helps reduce the risk of stroke and systemic embolism in appropriately selected patients.

Indications:

Warwise 1 Tablet is indicated for the prevention and treatment of thromboembolic disorders, including:

- **Prevention of Stroke and Systemic Embolism in Atrial Fibrillation**
- **Treatment and Prevention of Venous Thromboembolism (DVT and PE)**
- **Prevention of Thromboembolism associated with Mechanical Heart Valves**
- Other thromboembolic conditions as determined by the physician

---

Dosage and Administration:

- **Recommended Dose:** The dose is **individualized according to INR response** and the patient's clinical condition. There is no fixed routine dose for all patients.
- Take once daily at approximately the **same time each day**.
- Swallow the tablet with water, with or without food.
- Regular **PT/INR monitoring is essential** during treatment.

- Never change the dose or discontinue treatment without medical advice.
- Maintain a reasonably consistent intake of **vitamin K-containing foods** rather than making sudden major dietary changes.

Precautions:

- **INR monitoring is mandatory** to maintain the desired anticoagulant effect and reduce bleeding risk.
- Report unusual bruising, prolonged bleeding, blood in urine or stools, vomiting blood, severe headache, or unexplained weakness immediately.
- Inform the physician before any **surgery, dental procedure, or invasive procedure**.
- Use cautiously in patients with liver disease, uncontrolled hypertension, bleeding disorders, or increased risk of falls/trauma.
- Avoid excessive alcohol consumption.
- Patients should inform healthcare professionals that they are taking warfarin.
- **Pregnancy:** Warfarin is generally contraindicated during pregnancy except in specific specialist-managed circumstances because it can cause fetal harm.
- **Lactation:** Warfarin is generally considered compatible with breastfeeding when clinically indicated.

Contraindications:

- Active clinically significant bleeding.
- Known hypersensitivity to warfarin or any component.
- Conditions associated with a high risk of serious bleeding.
- Hemorrhagic disorders.
- Certain recent or planned surgeries involving the central nervous system or eye, depending on clinical circumstances.
- **Pregnancy**, except in selected specialist-managed situations where benefits outweigh risks.
- Inability to safely comply with required anticoagulation monitoring.

Adverse Reactions:

Common:

- Easy bruising
- Nosebleed
- Gum bleeding
- Prolonged bleeding from minor cuts
- Anemia

Less Common but Serious:

- **Major or life-threatening bleeding**
- Gastrointestinal bleeding
- Intracranial hemorrhage
- Hematuria
- Skin necrosis
- Purple toe syndrome
- Severe hypersensitivity reactions

Patients should seek **immediate medical attention** for uncontrolled bleeding, black/tarry stools, blood in vomit, severe headache, sudden weakness, or loss of consciousness.

Drug Interactions:

Warfarin has numerous clinically important interactions.

- **Antiplatelet agents and NSAIDs** may significantly increase bleeding risk.
- Many **antibiotics and antifungal medicines** can increase or decrease the anticoagulant effect.
- Other anticoagulants may produce additive anticoagulation.
- Medicines that induce or inhibit hepatic drug metabolism may substantially alter INR.
- **Herbal products and supplements**, including preparations containing St. John's wort, may affect warfarin activity.
- Sudden changes in dietary **vitamin K** intake can alter INR.
- Alcohol intake can affect anticoagulation and should be discussed with the physician.
- Patients should consult their physician or pharmacist before starting or stopping **any new medicine, OTC product, or supplement**.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture and direct sunlight.
- Keep tablets in the original packaging.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. **Warwise 1 Tablet must be used under the supervision of a registered medical practitioner with regular PT/INR monitoring.** The appropriate dose varies substantially between patients and depends on the indication, target INR, age, diet, liver function, genetics, and interacting medicines.

Important: Warfarin has a narrow therapeutic range and numerous drug and food interactions. Patients should **never adjust, stop, or restart warfarin on their own.**

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for