Information For the Use of
Warfarin (2mg) Tablet

Warwise 2 Tablet

Warwise 2 Tablet

Composition:

Each tablet contains:

- **Warfarin** – 2 mg

Pharmacological Action:

Warfarin is an **oral anticoagulant** that reduces the production of vitamin K-dependent clotting factors.

- **Vitamin K Antagonism:** Inhibits vitamin K epoxide reductase, reducing the regeneration of active vitamin K.
- **Reduced Clotting Factors:** Decreases synthesis of Factors II, VII, IX, and X, along with anticoagulant proteins C and S.
- **Anticoagulant Action:** Prolongs clotting time and helps prevent the formation and growth of blood clots.
- **Thromboembolic Protection:** Helps reduce the risk of stroke and systemic embolism in appropriately selected patients.

Indications:

Warwise 2 Tablet is indicated for the prevention and treatment of thromboembolic disorders, including:

- **Prevention of Stroke and Systemic Embolism in Atrial Fibrillation**
- **Treatment and Prevention of Deep Vein Thrombosis (DVT)**
- **Treatment and Prevention of Pulmonary Embolism (PE)**
- **Prevention of Thromboembolism associated with Mechanical Heart Valves**
- Other thromboembolic conditions as determined by the physician

---

Dosage and Administration:

- **Recommended Dose:** The dose is individualized according to the patient's **PT/INR** and clinical indication. There is no single standard dose suitable for every patient.
- Take once daily at approximately the **same time each day**.
- Swallow the tablet with water, with or without food.

- Regular **PT/INR monitoring is essential**.
- Never change the dose, skip doses, or discontinue treatment without medical advice.
- Maintain a reasonably consistent intake of **vitamin K-containing foods** and avoid sudden major dietary changes.

Precautions:

- Regular **INR monitoring** is required to maintain effective and safe anticoagulation.
- Report unusual bruising, prolonged bleeding, blood in urine or stools, vomiting blood, severe headache, or unexplained weakness immediately.
- Inform the physician before any **surgery, dental procedure, or invasive procedure**.
- Use cautiously in patients with liver disease, uncontrolled hypertension, bleeding disorders, or increased risk of falls or trauma.
- Avoid excessive alcohol consumption.
- Always inform healthcare professionals that you are taking warfarin.
- **Pregnancy:** Warfarin can cause fetal harm and is generally contraindicated during pregnancy except in selected specialist-managed circumstances.
- **Lactation:** Warfarin is generally considered compatible with breastfeeding when clinically indicated.

Contraindications:

- Active clinically significant bleeding.
- Known hypersensitivity to warfarin or any component of the formulation.
- Conditions associated with a high risk of serious bleeding.
- Hemorrhagic disorders.
- Certain recent or planned surgeries involving the central nervous system or eye, depending on clinical circumstances.
- **Pregnancy**, except in selected specialist-managed situations.
- Inability to safely comply with required anticoagulation monitoring.

Adverse Reactions:

Common:

- Easy bruising
- Nosebleed
- Gum bleeding
- Prolonged bleeding from minor cuts
- Anemia

Less Common but Serious:

- **Major or life-threatening bleeding**
- Gastrointestinal bleeding
- Intracranial hemorrhage
- Hematuria
- Skin necrosis
- Purple toe syndrome
- Severe hypersensitivity reactions

Seek immediate medical attention for uncontrolled bleeding, black/tarry stools, blood in vomit, severe headache, sudden weakness, or loss of consciousness.

Drug Interactions:

Warfarin has numerous clinically significant drug and food interactions.

- **Antiplatelet agents and NSAIDs** may increase bleeding risk.
- Many **antibiotics and antifungal medicines** can increase or decrease the anticoagulant effect.
- Other anticoagulants may produce additive anticoagulation.
- Medicines that inhibit or induce hepatic drug metabolism can substantially alter INR.
- **Herbal products and supplements**, including St. John's wort, may affect warfarin activity.
- Sudden changes in dietary **vitamin K** intake can alter INR.
- Alcohol consumption can affect anticoagulation.
- Consult a physician or pharmacist before starting or stopping any prescription medicine, OTC product, or supplement.

Storage:

- Store in a cool, dry place below **25°C**.
- Protect from moisture and direct sunlight.
- Keep tablets in the original packaging.
- Keep out of reach of children.

Note:

This information is intended as a general prescribing guide. **Warwise 2 Tablet must be used under the supervision of a registered medical practitioner with regular PT/INR monitoring.** The appropriate warfarin dose varies considerably between patients and depends on the indication, target INR, age, diet, liver function, and interacting medicines.

Important: Warfarin has a narrow therapeutic range. Patients should **never adjust, stop, or restart Warwise 2 Tablet on their own.**

Manufactured in India for:

Cafoli Lifecare Pvt. Ltd.
(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113
Corporate Office :
Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113
TM: Trademark Applied for