

The (normal) simple linear regression model:

$$Y_i = \beta_0 + \beta_1 x_i + \varepsilon_i,$$

where ε_i 's are independent $\text{Normal}(0, \sigma^2)$ (iid $\text{Normal}(0, \sigma^2)$).

β_0 , β_1 , and σ^2 are unknown model parameters.

$$\left[\begin{array}{l} SXX = \sum (x_i - \bar{x})^2 = \sum x_i^2 - \frac{(\sum x_i)^2}{n} \\ SXY = \sum (x_i - \bar{x})(y_i - \bar{y}) = \sum (x_i - \bar{x})y_i = \sum x_i y_i - \frac{(\sum x_i)(\sum y_i)}{n} \\ SY Y = \sum (y_i - \bar{y})^2 = \sum y_i^2 - \frac{(\sum y_i)^2}{n} \\ \text{Slope} \quad \hat{\beta}_1 = \frac{SXY}{SXX} \quad \text{Y-intercept} \quad \hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x} \end{array} \right]$$

Suppose x_i 's are fixed (not random).

$\Rightarrow Y_i$'s are independent $\text{Normal}(\beta_0 + \beta_1 x_i, \sigma^2)$ random variables.

$$\hat{\beta}_1 = \frac{\sum (x_i - \bar{x}) Y_i}{\sum (x_i - \bar{x})^2} \sim N \left(\beta_1, \frac{\sigma^2}{\sum (x_i - \bar{x})^2} \right)$$

$$\hat{\beta}_0 = \bar{Y} - \hat{\beta}_1 \bar{x} \sim N \left(\beta_0, \frac{\sigma^2 \sum x_i^2}{n \sum (x_i - \bar{x})^2} \right) = N \left(\beta_0, \sigma^2 \left(\frac{1}{n} + \frac{\bar{x}^2}{\sum (x_i - \bar{x})^2} \right) \right)$$

$$S_e^2 = \frac{RSS}{n-2} = \frac{1}{n-2} \sum (Y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2 \quad \frac{RSS}{\sigma^2} = \frac{(n-2) S_e^2}{\sigma^2} \sim \chi^2(n-2)$$

1. The owner of *Momma Leona's Pizza* restaurant chain believes that if a restaurant is located near a college campus, then there is a linear relationship between sales and the size of the student population. Suppose data were collected from a sample of 10 *Momma Leona's Pizza* restaurants located near college campuses. For the i th restaurant in the sample, x_i is the size of the student population (in thousands) and y_i is the quarterly sales (in thousands of dollars). The values of x_i and y_i for the 10 restaurants in the sample are summarized in the following table:

Restaurant	Student Population (1000s)	Quarterly Sales (\$1000s)
i	x_i	y_i
1	2	58
2	6	105
3	8	88
4	8	118
5	12	117
6	16	137
7	20	157
8	20	169
9	22	149
10	26	202

$$\bar{x} = 14, \quad \bar{y} = 130$$

$$SXX = 568$$

$$SXY = 2,840$$

$$SYY = 15,730$$


$$\hat{\beta}_1 = 5, \quad \hat{\beta}_0 = 60$$

$$\hat{y} = 60 + 5 \cdot x$$

$$RSS = 1,530$$

$$R^2 = 0.9027$$

$$s_e^2 = 191.25$$



Confidence interval for β_1 : $\hat{\beta}_1 \pm t_{\alpha/2} \cdot \frac{s_e}{\sqrt{\sum (x_i - \bar{x})^2}}$ $\hat{\beta}_1 \pm t_{\alpha/2} \cdot \frac{s_e}{\sqrt{SXX}}$

where $t_{\alpha/2}$ is the appropriate value of t-distribution

with $n - 2$ degrees of freedom.

Test statistic for $H_0: \beta_1 = \beta_{10}$:

$$T = \frac{\hat{\beta}_1 - \beta_{10}}{\frac{s_e}{\sqrt{\sum (x_i - \bar{x})^2}}} = \frac{\hat{\beta}_1 - \beta_{10}}{\frac{s_e}{\sqrt{SXX}}} \quad (n - 2 \text{ degrees of freedom})$$

a) Construct a 90% confidence interval for β_1 .

$$\hat{\beta}_1 \pm t_{\alpha/2} \cdot \frac{s_e}{\sqrt{SXX}} \quad 10 - 2 = 8 \text{ degrees of freedom,} \quad t_{0.05} = 1.860.$$

$$5 \pm 1.860 \cdot \frac{13.83}{\sqrt{568}} \quad \mathbf{5 \pm 1.08} \quad \mathbf{(3.92, 6.08)}$$

b) Test the assumption that students do not affect the sales. That is, test $H_0: \beta_1 = 0$ vs. $H_1: \beta_1 \neq 0$ (the significance of regression test).
Use $\alpha = 0.01$.

Test Statistic:

$$T = \frac{\hat{\beta}_1 - \beta_{10}}{\frac{s_e}{\sqrt{SXX}}} = \frac{5 - 0}{\frac{\sqrt{191.25}}{\sqrt{568}}} = 8.616.$$

Rejection Region:

Reject H_0 if $T < -t_{0.005}(10 - 2 = 8 \text{ df})$ or $T > t_{0.005}(8 \text{ df})$

$$\pm t_{0.005}(8 \text{ df}) = \pm 3.355.$$

Reject H_0

c) That is, test $H_0: \beta_1 = 4$ vs. $H_1: \beta_1 > 4$. Use $\alpha = 0.05$.

Test Statistic:

$$T = \frac{\hat{\beta}_1 - \beta_{10}}{s_e / \sqrt{SXX}} = \frac{5 - 4}{\sqrt{191.25} / \sqrt{568}} = 1.723.$$

Rejection Region:

Reject H_0 if $T > t_{0.05}(8 \text{ df})$

$$t_{0.05}(8 \text{ df}) = 1.860.$$

Do NOT Reject H_0

($0.05 < \text{p-value} < 0.10$)

- - - - -

Confidence interval for β_0 :

$$\hat{\beta}_0 \pm t_{\alpha/2} \cdot s_e \sqrt{\frac{1}{n} + \frac{\bar{x}^2}{SXX}}$$

where $t_{\alpha/2}$ is the appropriate value of t-distribution

with $n - 2$ degrees of freedom.

Test statistic for $H_0: \beta_0 = \beta_{00}$:

$$T = \frac{\hat{\beta}_0 - \beta_{00}}{s_e \sqrt{\frac{1}{n} + \frac{\bar{x}^2}{SXX}}} \quad (n - 2 \text{ degrees of freedom})$$

d) Construct a 90% confidence interval for β_0 .

$$60 \pm 1.860 \cdot \sqrt{191.25} \cdot \sqrt{\frac{1}{10} + \frac{14^2}{568}} \quad \quad \quad \mathbf{60 \pm 17.16}$$

e) Test $H_0: \beta_0 = 75$ vs. $H_1: \beta_0 < 75$. Use a 5% level of significance.

$$\text{Test Statistic: } T = \frac{60 - 75}{\sqrt{191.25} \sqrt{\frac{1}{10} + \frac{14^2}{568}}} = -1.626.$$

Rejection Region: Reject H_0 if $T < -t_{0.05}(8 \text{ df}) = -1.860$.

Do NOT Reject H_0

Confidence interval for σ^2 :

$$\left(\frac{(n-2)s_e^2}{\chi_{\alpha/2}^2}, \frac{(n-2)s_e^2}{\chi_{1-\alpha/2}^2} \right) \quad \left(\frac{n\hat{\sigma}^2}{\chi_{\alpha/2}^2}, \frac{n\hat{\sigma}^2}{\chi_{1-\alpha/2}^2} \right)$$

where $\chi_{1-\alpha/2}^2$ and $\chi_{\alpha/2}^2$ are the appropriate values of χ^2 distribution
with $n - 2$ degrees of freedom.

f) Construct a 95% confidence interval for σ^2 .

$$\chi_{0.025}^2(8 \text{ df}) = 17.54, \quad \chi_{0.975}^2(8 \text{ df}) = 2.180.$$

$$\left(\frac{8 \cdot 191.25}{17.54}, \frac{8 \cdot 191.25}{2.180} \right) \quad (87.229, 701.835)$$

Mean response (y) for a fixed value of x : $\mu(x) = \mu_{y|x} = \beta_0 + \beta_1 x$.

To estimate $\mu(x)$, use $\hat{y} = \hat{\beta}_0 + \hat{\beta}_1 x$.

$$E(\hat{Y}) = \mu(x) = \beta_0 + \beta_1 x. \quad \text{Var}(\hat{Y}) = \sigma^2 \left(\frac{1}{n} + \frac{(x - \bar{x})^2}{SXX} \right).$$

Confidence interval for $\mu(x)$:

$$\hat{y} \pm t_{\alpha/2} \cdot s_e \sqrt{\frac{1}{n} + \frac{(x - \bar{x})^2}{SXX}}$$

where $t_{\alpha/2}$ is the appropriate value of t-distribution
with $n - 2$ degrees of freedom.

Prediction interval for a future value of y corresponding to a given value of x :

$$\hat{y} \pm t_{\alpha/2} \cdot s_e \sqrt{1 + \frac{1}{n} + \frac{(x - \bar{x})^2}{SXX}} \quad (\text{limits of prediction})$$

where $t_{\alpha/2}$ is the appropriate value of t-distribution with $n - 2$ degrees of freedom.

g) Construct a 95% confidence interval for $\mu(x = 10)$.

$$110 \pm 2.306 \cdot \sqrt{191.25} \cdot \sqrt{\frac{1}{10} + \frac{(10 - 14)^2}{568}}$$

$$\mathbf{110 \pm 11.42}$$

h) Construct a 95% confidence interval for $\mu(x = 38)$.


$$250 \pm 2.306 \cdot \sqrt{191.25} \cdot \sqrt{\frac{1}{10} + \frac{(38 - 14)^2}{568}}$$

$$\mathbf{250 \pm 33.66}$$

i) Construct a 95% prediction interval for a future value of y corresponding to $x = 38$.
(Construct 95% limits of prediction if $x = 38$.)

$$250 \pm 2.306 \cdot \sqrt{191.25} \cdot \sqrt{1 + \frac{1}{10} + \frac{(38 - 14)^2}{568}}$$

$$\mathbf{250 \pm 46.37}$$



- j) University of Illinois at Urbana-Champaign has 38 thousand students. The owner of *Momma Leona's Pizza* restaurant chain would agree to open a restaurant near the UIUC campus, but only if there is enough evidence that the average quarterly sales would be over \$225,000. Test $H_0: \mu(x=38) = 225$ vs. $H_1: \mu(x=38) > 225$. Use $\alpha = 0.05$.

Test Statistic:
$$T = \frac{250 - 225}{\sqrt{191.25} \sqrt{\frac{1}{10} + \frac{(38-14)^2}{568}}} = 1.713.$$


Rejection Region: $\text{Reject } H_0 \text{ if } T > t_{0.05}(8 \text{ df}) = 1.860.$

Do NOT Reject H_0

Note that $\beta_0 = \mu(x=0).$

$$\hat{\beta}_0 = \hat{y} \quad \text{if } x = 0.$$

- b) Test the assumption that students do not affect the sales. That is, test $H_0: \beta_1 = 0$ vs. $H_1: \beta_1 \neq 0$ (the significance of regression test).
Use $\alpha = 0.01$.



Here $V_0 = \{ a \mathbf{1}, a \in \mathbf{R} \}$, $\dim(V_0) = 1$, $\hat{\mathbf{Y}}_0 = [\bar{Y}, \bar{Y}, \dots, \bar{Y}]^T$,
 $V = \{ a_0 \mathbf{1} + a_1 \mathbf{x}, a_0, a_1 \in \mathbf{R} \}$, $\dim(V) = 2$.

$$\sum (y_i - \bar{y})^2 = \sum (y_i - \hat{y}_i)^2 + \sum (\hat{y}_i - \bar{y})^2$$

$$\text{Since } \hat{y}_i = \hat{\alpha} + \hat{\beta} x_i = (\bar{y} - \hat{\beta} \bar{x}) + \hat{\beta} x_i = \bar{y} + \hat{\beta} (x_i - \bar{x}),$$

$$\text{SSRegression} = \sum (\hat{y}_i - \bar{y})^2 = \sum \hat{\beta}^2 (x_i - \bar{x})^2 = \hat{\beta}^2 \sum (x_i - \bar{x})^2 = \hat{\beta}^2 \text{SXX}.$$

ANOVA table:

Source	SS	DF	MS	F
Regression	$\sum (\hat{y}_i - \bar{y})^2 = 14,200$	1	14,200	74.248366
Error	$\sum (y_i - \hat{y}_i)^2 = 1,530$	$n - 2 = 8$	191.25	
Total	$\sum (y_i - \bar{y})^2 = 15,730$	$n - 1 = 9$		

Rejection Region: Reject H_0 if $F > F_{0.01}(1, 8) = 11.26$.

Reject H_0