

# WS #18 - Support Vector Machines 3

Wednesday, November 12, 2025

Your Name: \_\_\_\_\_

Names of people you worked with: \_\_\_\_\_

What was the number one reason you came to the Claremont Colleges? As such, have you been pleased or disappointed?

## Task:

Consider two 2-dimensional data points:  $\mathbf{A} = (1, 2)$ ,  $\mathbf{B} = (2, 4)$ . Use the following function to map the two data points into the indicated six-dimensional space:

$$\phi(x) = (x_1^2, x_2^2, \sqrt{2}x_1x_2, \sqrt{2}x_1, \sqrt{2}x_2, 1)$$

Do the following computations:

a.  $\phi(\mathbf{A}) =$

b.  $\phi(\mathbf{B}) =$

c.  $\phi(\mathbf{A}) \cdot \phi(\mathbf{B}) = \phi(\mathbf{A})^t \phi(\mathbf{B}) =$

The kernel function corresponding to the projection above is,

$$K(\mathbf{x}, \mathbf{y}) = (\mathbf{x}^t \mathbf{y} + 1)^2$$

(A degree-two polynomial kernel. More generally, we could choose a polynomial kernel of arbitrary degree:  $K(\mathbf{x}, \mathbf{y}) = (\mathbf{x}^t \mathbf{y} + \text{const})^d$ .)

d.  $K(\mathbf{A}, \mathbf{B}) =$

**Solution:**

a.  $\phi(\mathbf{A}) = (1, 4, 2\sqrt{2}, \sqrt{2}, 2\sqrt{2}, 1)$

b.  $\phi(\mathbf{B}) = (4, 16, 8\sqrt{2}, 2\sqrt{2}, 4\sqrt{2}, 1)$

c.  $\phi(\mathbf{A}) \cdot \phi(\mathbf{B}) = \phi(\mathbf{A})^t \phi(\mathbf{B}) = (4 + 64 + 16 \cdot 2 + 2 \cdot 2 + 8 \cdot 2 + 1) = 121$

$$K(\mathbf{x}, \mathbf{y}) = (\mathbf{x}^t \mathbf{y} + 1)^2$$

d.  $K(\mathbf{A}, \mathbf{B}) = ((1 \cdot 2 + 2 \cdot 4) + 1)^2 = 11^2 = 121$