

10.20 The following data give the number of casualties for 103 suicide attacks in Israel with explosives over a three-year period from November 6, 2000 to November 3, 2003 when there was a steep drop (the early period of the first “Intifada”). These data are provided by the International Policy Institute for Counter-Terrorism, and subsetting by Harrison (2006), and modeled in Kyung *et al.* (2011).

0	3	81	38	29	126	6	10	1	1
67	50	3	27	0	2	0	63	15	58
57	0	0	0	0	123	4	71	71	20
17	65	4	49	5	35	57	71	0	12
67	59	5	52	62	0	75	0	0	106
30	0	3	45	4	31	32	180	0	1
91	49	61	51	3	0	1	9	0	2
151	26	8	8	75	199	12	2	2	1
93	0	13	21	145	0	0	13	0	2
141	2	65	0	105	0	61	6	27	53
20	5	0							

There are two modes, with the larger one at zero. Fit a two-component gamma mixture model using data augmentation for p (the probability of being in the right-mode) with a beta distribution, and with a binary assignment vector as part of the Gibbs sampler. At iteration t the steps are:

- ▷ Draw p from $\mathcal{BE}(\alpha + n_1, \beta + n_2)$.
- ▷ For each case i , draw $I_i \sim \mathcal{BR}(p)$.
- ▷ If $I_i = 0$, draw from $\mathcal{G}(\alpha_0, \beta_0)$.
- ▷ If $I_i = 1$, draw from $\mathcal{G}(\alpha_1, \beta_1)$.
- ▷ Determine n_1 and n_2 from the I_i assignments.