UNIT 4
Introduction to Data Science

Model Development: Simple and Multiple Regression – Model Evaluation using Visualization – Residual Plot –Distribution Plot – Polynomial Regression and Pipelines – Measures for In-sampleEvaluation – Prediction and Decision Making.

Model Development

Regression analysis is a predictive modelling technique that assesses the relationship between dependent (i.e., the goal/target variable) and independent factors.

Forecasting, time series modelling, determining the relationship between variables, and predicting continuous values can all be done using regression analysis. Just to give you an analogy, Regression is the best way to study the relationship between household areas and a driver's household electricity cost.

Dependent Variable - The dependent variable is essentially the "outcome" you're trying to understand or predict.

Independent Variable - Independent variables are the "factors" that might influence or cause changes in the dependent variable.

SIMPLE AND MULTIPLE REGRESSION :

- Simple and multiple regression are **supervised machine learning methods**.
- Used to predict a continuous numerical output (Y) based on input features (X).
- **Types :**
    1. **Simple regression** uses **one independent variable** to predict a single dependent outcome using a straight line.
    2. **Multiple regression** uses **two or more independent variables** to predict that same outcome, fitting a multi-dimensional plane to capture how several combined factors affect the target value

DIFFERENCE BETWEEN SIMPLE AND MULTIPLE REGRESSION

Simple Regression	Multiple Regression
Simple uses one X variable	Multiple uses two or more (X_1, X_2, etc.).
Simple regression draws a 2D straight line	Multiple regression fits a multi-dimensional space or plane
Simple looks like Y = a + bX	Multiple expands to \(Y = a + b_1X_1 + b_2X_2 + ... + b_nX_n\).
Simple is easier to calculate but often too basic	Multiple is realistic because outcomes usually depend on many combined forces

Terminologies Related to the Regression Analysis:

- **Dependent Variable: The main factor in Regression analysis which we want to predict or understand is called the dependent variable. It is also called target variable.**

- **Independent Variable: The factors which affect the dependent variables or which are used to predict the values of the dependent variables are called independent variables, also called as a predictor.**


- **Outliers: Outlier is an observation which contains either very low value or very high value in comparison to other observed values. An outlier may hamper the result, so it should be avoided.**

- **Underfitting and Overfitting: If our algorithm works well with the training dataset but not well with test dataset, then such problem is called Overfitting. And if our algorithm does not perform well even with training dataset, then such problem is called underfitting.**

LINEAR REGRESSION

- **Linear regression is a statistical regression method which is used for predictive analysis.**
- **It is one of the very simple and easy algorithms which works on regression and shows the relationship between the continuous variables.**
- **It is used for solving the regression problem in machine learning.**
- **Linear regression shows the linear relationship between the independent variable (X-axis) and the dependent variable (Y-axis), hence called linear regression.**
- **If there is only one input variable (x), then such linear regression is called simple linear regression. And if there is more than one input variable, then such linear regression is called multiple linear regression.**

The relationship between variables in the linear regression model can be explained using the below image. Here predicting the salary of an employee on the basis of the year of experience.



- Below is the mathematical equation for Linear regression:

$$Y = aX + b$$

Here, Y = dependent variables (target variables),
X= Independent variables (predictor variables),
a and b are the linear coefficients

Some popular applications of linear regression are:

- **Analyzing trends and sales estimates**
- **Salary forecasting**
- **Real estate prediction**
- **Arriving at ETAs in traffic.**

Multiple Regression :

- **Multiple linear regression is used to estimate the relationship between two or more independent variables and one dependent variable. You can use multiple linear regression when you want to know:**
  1. **How strong the relationship is between two or more independent variables and one dependent variable (e.g. how rainfall, temperature, and amount of fertilizer added affect crop growth).**
  2. **The value of the dependent variable at a certain value of the independent variables (e.g. the expected yield of a crop at certain levels of rainfall, temperature, and fertilizer addition).**

Multiple linear regression example.

- **You are a public health researcher interested in social factors that influence heart disease. You survey 500 towns and gather data on the percentage of people in each town who smoke, the percentage of people in each town who bike to work, and the percentage of people in each town who have heart disease.**
- **Because you have two independent variables and one dependent variable, and all your variables are quantitative, you can use multiple linear regression to analyze the relationship between them.**

How to perform a multiple linear regression

Multiple linear regression formula

The formula for a multiple linear regression is:

$$y = \beta_0 + \beta_1 X_1 + \ldots + \beta_n X_n + \epsilon$$

- y = the predicted value of the dependent variable
- B_0 = the y-intercept (value of y when all other parameters are set to 0)
- $B_1 X_1$ = the regression coefficient (B_1) of the first independent variable (X_1) (a.k.a. the effect that increasing the value of the independent variable has on the predicted y value)
- … = do the same for however many independent variables you are testing
- $B_n X_n$ = the regression coefficient of the last independent variable
- ϵ = model error (a.k.a. how much variation there is in our estimate of y)

Simple Linear Regression: **Simple linear regression is used when a single independent variable predicts a dependent variable**

- **The association between two variables is established using a straight line in Simple Linear Regression.**

- **It tries to create a line that is as near to the data as possible by determining the slope and intercept, which define the line and reduce regression errors.**
- **There is a single x and y variable**

Example:

Suppose we want to understand the relationship between a company's stock price (dependent variable) and the company's quarterly earnings (independent variable). For several quarters, we collect historical data on the company's earnings and stock prices. And by performing simple linear regression, we can identify the linear relationship between earnings and stock prices


Multiple Linear Regression:

Multiple regression extends linear regression by considering multiple independent variables to predict the dependent variable.

- **Multiple linear regressions are based on the presumption that both the dependent and independent variables, or Predictor and Target variables, have a linear relationship.**
- **There are two types of multilinear regressions: linear and nonlinear.**
- **It has one or more x variables and one or more y variables, or one dependent variable and two or more independent variables**

Example:

In real estate, we can predict the selling price of a house based on various factors such as area, number of bedrooms, number of floors, and location. This is where multiple linear regression comes into play.



Types of Regression Analysis:

Simple Linear Regression

Simple linear regression is used to model the relationship between two variables, where one is considered the independent variable (predictor) and the other is the dependent variable (outcome).

Multiple Linear Regression

Multiple linear regression extends simple linear regression to model relationships between multiple independent variables and a single dependent variable.

Logistic Regression

Logistic regression is used when the dependent variable is binary (two possible outcomes). It models the probability of a particular outcome occurring.

Polynomial Regression

Polynomial regression is used when the relationship between the independent and dependent variables follows a polynomial curve and is not linear.

Non-linear Regression

Non-linear regression is used when the relationship between the dependent and independent variables can take various functional forms.

Model Evaluation using Visualization

- **Model evaluation is the process of utilizing various evaluation metrics to enhance and assess a model's performance.**
- **It prevents overfitting, enhances accuracy, and ensures models achieve goals efficiently and effectively.**

Residual Plot

Residuals:

Residual analysis is a powerful statistical technique used to assess the accuracy of regression models.

- **A residual is the difference between a variable's observed value and the variable's predicted value based on a statistical or ML model.**
- **Researchers and analysts need this technique to make better decisions about the validity and reliability of their statistical models.**

Residual analysis


- **Residuals are used to assess the validity of a statistical model.**
- **The model is considered a good fit if the residuals are randomly distributed.**
- **If there are patterns in the residuals, then the model is not accurately capturing the relationship between the variables.**

- **It may need to be improved, or another model may need to be selected.**

Residual Plot

Residual plots are graphical representations of the residuals against the predictor variables in a regression analysis. These plots help assess the assumptions and adequacy of the regression model.

- **A residual plot is a scatterplot in which the X-axis represents the independent or target variable, and the Y-axis represents residual values based on the ML model.**
- **A residual plot is used to identify the underlying patterns in the residual values.**




- A **residual plot** is a graph that shows the differences (residuals) between actual data values and the predicted values from a regression model

Characteristics of Good Residual Plots

A few characteristics of a good residual plot are as follows:

1. **It has a high density of points close to the origin and a low density of points away from the origin**
2. **It is symmetric about the origin**


To explain why below fig is a good residual plot based on the characteristics above, we project all the residuals onto the y-axis. As seen in Figure 3b, we end up with a normally distributed curve; satisfying the assumption of the normality of the residuals.




Finally, one other reason this is a good residual plot is, that independent of the value of an independent variable (x-axis), the residual errors are approximately distributed in the same manner. In other words, we do not see any patterns in the value of the residuals as we move along the x-axis.

Hence, this satisfies our earlier assumption that regression model residuals are independent and normally distributed.

Using the characteristics of the bad residual plot: this plot has high density far away from the origin and low density close to the origin. Also, when we project the residuals on the y-axis, we can see the distribution curve is not normal.



Example of Bad Residual plot



Project onto the y-axis

Distribution plots

Distribution plots visualize how data points are spread across a range of values, allowing for comparison of ranges and identification of patterns, outliers, and concentration of data

Common types include histograms, which show the frequency of data in bins, and density plots, which provide a smoothed curve estimate of the distribution. Other forms are box plots, summarizing quartiles and outliers, and violin plots, offering a more detailed view than box plots.

Purpose and Use

- **Data Exploration: To understand the shape and spread of data.**
- **Comparison: To compare the distribution of different groups of data.**
- **Outlier Detection: To identify extreme values that deviate from the main data cluster.**
- **Assumption Checking: To visually assess if data conforms to a theoretical distribution (e.g., normal distribution).**


Common Types of Distribution Plots

- **Histograms: Divide data into bins (intervals) and show the frequency of data points within each bin using bars.**
- **Box Plots: Summarize the data's distribution by displaying the median, quartiles, and potential outliers.**
- **Violin Plots: Combine aspects of box plots with a kernel density estimation to show both the summary statistics and the probability density of the data.**
- **Kernel Density Estimation (KDE) Plot: Smooths the data to create a continuous curve, showing the probability density of the data at different values.**
- **Cumulative Distribution Plots (CDF Plots): Display the cumulative frequency of data, showing the proportion of data that falls below a specific value.**
- **Quantile-Quantile (Q-Q) Plots: Compare the empirical distribution of sample data against the theoretical values expected from a given distribution to assess if the sample data comes from that distribution.**

Examples:


Histogram: Uses adjacent vertical bars to show the frequency of continuous data divided into equal intervals (bins).

- *Example*: Plotting the distribution of student test scores (e.g., binning scores into 60-70, 70-80) to see how many students fall into each range.




Box Plot (Box-and-Whisker Plot): Summarizes data spread using the median, quartiles, and potential outliers via a central box and outer "whiskers."

- *Example*: Comparing annual salaries across different departments to spot outliers or minimum/maximum earnings.




Density Plot (KDE Plot): Displays a smooth, continuous curve representing the probability density of the data instead of discrete bars.

- *Example*: Visualizing the smooth distribution of customer ages in a retail dataset




Violin Plot: Combines a Box Plot summary with a smooth Kernel Density Estimation curve on both sides.
A violin plot combines the features of a box plot with a kernel density plot.

- *Example*: Showing both summary statistics and multi-peaked (multimodal) distributions of daily temperature readings.



Polynomial Regression:

- **Polynomial Regression is a form of regression analysis in which the relationship between the independent variables and dependent variables are modeled in the nth degree polynomial.**
- **Polynomial Regression models are usually fit with the method of least squares.*The least square method minimizes the variance of the coefficients.***
- **Polynomial Regression is a special case of Linear Regression where we fit the polynomial equation on the data with a curvilinear relationship between the dependent and independent variables.**
  - **Linear regression, which assumes a straight-line relationship.**
  - **Polynomial regression captures curved patterns by adding higher-degree terms of the independent variable (like x^2, x^3, etc.) to the model.**



In the simulated data above, the predictor variable on the x-axis is not linearly related to the outcome variable on the y-axis.


Instead, the relationship between these variables is better described by a curve. If we fit a linear regression model to these data, the model equation would be as follows:

$$Y \sim a + bX + \epsilon$$

where *a* is the intercept (the value at which the regression line cuts through the y-axis), *b* is the coefficient, and ϵ is an error term.

As seen in the plot above, this straight-line equation doesn't do a good job of capturing the non-linear relationship in the data. To address this, we can fit a polynomial regression model. This model is an extension of the previous one, but *X* is now added again as a second-degree polynomial. The equation for this model is

$$Y \sim a + bX + bX^2 + \epsilon$$



It's clear from a quick visual inspection that the polynomial model gives a closer fit to the curved data. This will lead to more accurate predictions of new values in test data.

More polynomial components can be added to the regression equation to fit more complex non-linear relationships. These would make the regression equation take this form:

$$Y \sim a + bX + bX^2 + bX^3 + \ldots + bX^n + \epsilon$$

Example Applications

- **Housing Price Prediction: To understand how factors like house size, age, and location non-linearly affect prices.**
- **Sales Forecasting: To capture complex growth patterns in sales over time.**
- **Agricultural Yields: To model the non-linear effects of temperature and rainfall on crop yields**

Types of Polynomial Regression

A quadratic equation is a general term for a second-degree polynomial equation. This degree, on the other hand, can go up to nth values. Polynomial regression can so be categorized as follows:

1. Linear – if degree as 1

2. Quadratic – if degree as 2

3. Cubic – if degree as 3 and goes on, on the basis of degree.

Polynomials	Form	Degree	Examples
Linear Polynomial	$p(x): ax+b, a \neq 0$	Polynomial with Degree 1	$x + 8$
Quadratic Polynomial	$p(x): ax^2+b+c, a \neq 0$	Polynomial with Degree 2	$3x^2-4x+7$
Cubic Polynomial	$p(x): ax^3+bx^2+cx, a \neq 0$	Polynomial with Degree 3	$2x^3+3x^2+4x+6$

Data science pipeline

- **A Data Science Pipeline is a collection of processes that transform raw data into actionable business answers.**

- **The Data Science Pipeline refers to the process and tools used to collect raw data from various sources, analyze it, and present the results in a Comprehensible Format.**

- **Companies use the process to answer specific business questions and generate actionable insights from real-world data. To find this information, all available Datasets, both External and Internal, are analyzed.**

It involves several stages, including data collection, data preprocessing (cleaning and transforming), exploratory data analysis (EDA), feature engineering, model development, model evaluation, and finally, deployment and monitoring.

This structured process ensures data quality, reproducibility, and scalability, streamlining the entire process from raw data to meaningful results

1. **Data collection**

   Data is collected from various sources, including data warehouses, APIs, business tools, or web scraping apps. When scraping location-specific content, the best residential proxies can help ensure accurate, geo-targeted data collection while avoiding IP bans.

2. **Data cleaning:**

   Incomplete, irrelevant, and unstructured data is cleaned and formatted properly so there are no issues with processes like exploratory data analysis later on. Missing values are added to unstructured data.

3. **Data transformation:**

Data is transformed into a format that is suitable for analysis later on. At this point, data goes through normalization, aggregation, or encoding categorical variables.

4. **Data exploration**

   Data distributions and relationships are explored, typically through data visualization such as graphs, dashboards, and reports. This allows even end users who are not familiar with data analytics to understand the relationships between data points.

5. **Feature engineering**

   New variables are created or existing ones are modified to improve the performance of the model and get essential information.

6. **Modeling**

   Machine learning algorithms are selected and trained to make predictions or classify data.

7. **Evaluating**

   Assessing how accurate the model is and how it performs based on metrics such as precision, recall.

8. **Deployment**

   The model is deployed in a real-life production environment where it can be used across apps and different platforms in various use cases.

9. **Monitoring and maintenance**
   The model performance is tracked continuously so you can make adjustments and ensure it stays accurate over time.

Through the automation and optimization of these processes, data-driven companies can speed up the time it takes to go from raw data to actionable insights.

Key Features of Data Science Pipelines

- **Continuous and Scalable Data Processing**
- **Cloud-based Elasticity and Agility.**
- **Data Processing Resources that are Self-Contained and Isolated.**
- **Access to a Large Amount of Data and the ability to self-serve.**
- **Disaster Recovery and High Availability.**

Purpose

Data science pipeline focuses on the entire process of developing and deploying machine learning models, from collecting data from various data sources, cleaning and evaluating it.

Data science pipelines are essential for several reasons:

Efficiency: Pipelines automate data processing tasks when possible, saving time and reducing manual effort.

Consistency: With a standardized pipeline, data undergoes the same process each time, minimizing inconsistencies.

Scalability: Pipelines allow data processing to scale, making it easier to handle large volumes of data or complex analyses.

Reproducibility: Pipelines create a repeatable process, making it easier to validate results or reproduce insights for future use.

Measures for In-sampleEvaluation

In-sample evaluation metrics assess a data science model's performance on the training data it was developed on, indicating how well it learned the training patterns.

Common in-sample metrics include Mean Squared Error (MSE) and Mean Absolute Error (MAE) for regression, and Accuracy, Precision, Recall, and the F1 Score for classification, alongside visualizations like Confusion Matrices and ROC Curves.

Regression Models

These metrics focus on the difference between predicted and actual numerical values.

- **Mean Squared Error (MSE): Measures the average squared difference between predicted and actual values. A lower MSE indicates a better fit.**
- **Mean Absolute Error (MAE): Calculates the average absolute difference between predicted and actual values. It's less sensitive to outliers than MSE.**
- **R-squared: Represents the proportion of the variance in the dependent variable that is predictable from the independent variables.**

MAE - Mean Absolute Error

Mean Absolute Error (MAE) measures the magnitude of the errors in a set of predictions, without considering their direction.

It is the average of the absolute differences between predictions and actual observations where all individual differences have equal weight.

$$MAE = \frac{1}{n}\sum_{i=1}^{n}|y_i - \hat{y}_i|$$

Where:

- n is the number of observations in the dataset.
- y_i represents the actual (observed) value for the i-th observation.
- $\hat{y}_i$ represents the predicted value for the i-th observation.
- $|\cdot|$ denotes the absolute value.

Lower values indicate better performance.

A value of 0 indicates no error. MAE = 0 means that the model is overfit.

It will work well on training data but not on testing data

$$\text{MAE} = \frac{1}{n}\sum_{i=1}^{n}|y_i - \hat{y}_i|$$

Step-by-step:

- Absolute errors: 2, 1, 2
- Add them: $2 + 1 + 2 = 5$
- Divide by 3:

$$\text{MAE} = \frac{5}{3} \approx \boxed{1.67}$$

MSE - Mean Squared Error

Mean Squared Error (MSE) measures the average of the squares of the errors i.e. the average squared difference between the estimated values and the actual values.

$$MSE = \frac{1}{n}\sum_{i=1}^{n}(y_i - \hat{y}_i)^2$$

Where:

- n is the number of observations in the dataset.
- y_i represents the actual (observed) value for the i-th observation.
- $\hat{y}_i$ represents the predicted value for the i-th observation.

Actual (y)	Predicted (ŷ)	Error (y - ŷ)	|Error|	Error²
10	12	-2	2	4

| 15 | 14 | 1 | 1 | 1 |
| 20 | 18 | 2 | 2 | 4 |

✅ Mean Squared Error (MSE)

Definition: MSE is the average of the squared differences between the actual values and the predicted values.

Formula:

$$\text{MSE} = \frac{1}{n}\sum_{i=1}^{n}(y_i - \hat{y}_i)^2$$

Step-by-step:

- Error^2 values: 4, 1, 4
- Add them: $4 + 1 + 4 = 9$
- Divide by 3 (number of data points):

$$\text{MSE} = \frac{9}{3} = \boxed{3.0}$$

Like MAE, lower values are better.

MSE is more sensitive to outliers than MAE because outliers skew the mean towards themselves.

When there are more outliers, whether the value of MSE would be greater or smaller depends on the dataset.

R-Squared - Coefficient of Determination

R-Squared represents the proportion of the variance for the dependent variable that is explained by the independent variables in a regression model.

$$R^2 = 1 - \frac{\sum_{i=1}^{n}(y_i - \hat{y}_i)^2}{\sum_{i=1}^{n}(y_i - \bar{y})^2}$$

Where:

- n is the number of observations.
- y_i is the actual (observed) value for the i-th observation.
- $\hat{y}_i$ is the predicted value for the i-th observation.
- $\bar{y}$ is the mean of the observed values y_i across all observations.

Formula:

$$R^2 = 1 - \frac{\text{Sum of Squared Errors (SSE)}}{\text{Total Sum of Squares (SST)}}$$

Metric	Value	Meaning
MSE	3.0	Average **squared** error; penalizes large errors more
MAE	1.67	Average **absolute** error; easier to understand
R^2	0.82	82% of variance in actual values is explained by the model

Prediction and Decision Making.

Prediction

Prediction is about using past data to forecast future events or outcomes. The goal is to build models that can generalize well to new, unseen data. In essence, prediction is a way of anticipating what will happen based on patterns in historical data.

Methods for Prediction:

1. **Supervised Learning: This is the most common form of predictive modeling. You train a model on labeled data (i.e., data with known outcomes), and the model learns to predict the output for new data.**
    - **Classification (predicting discrete labels): For example, predicting whether a customer will churn (Yes/No).**
    - **Regression (predicting continuous values): For example, predicting the price of a house based on its features (size, location, etc.).**
2. **Common algorithms:**
    - **Linear Regression**
    - **Logistic Regression**

- **Decision Trees**
- **Random Forests**
- **Support Vector Machines (SVM)**
- **Neural Networks (for more complex patterns)**

3. **Unsupervised Learning: Sometimes you don’t have labeled data, and you use unsupervised techniques to find patterns, clusters, or anomalies.**
    - **Clustering: Grouping similar items together, like segmenting customers based on purchasing behavior.**
    - **Dimensionality Reduction: Reducing the number of variables to simplify prediction or understand the data better (e.g., PCA - Principal Component Analysis).**
4. **Time Series Forecasting: In scenarios where you're predicting based on time-dependent data (e.g., stock prices, sales), models like ARIMA, LSTM (Long Short-Term Memory) networks, or Prophet are used to forecast future data points.**
5. **Ensemble Methods: To improve prediction accuracy, you can combine multiple models.**
    - **Bagging (e.g., Random Forest)**
    - **Boosting (e.g., XGBoost, LightGBM)**
6. **Deep Learning: For complex prediction tasks (e.g., image recognition, natural language processing), deep neural networks (e.g., Convolutional Neural Networks - CNNs or Recurrent Neural Networks - RNNs) can make highly accurate predictions, but they require large datasets.**

Key Considerations in Prediction:

- **Data Quality: The accuracy of predictions depends heavily on the quality and quantity of data.**

- **Feature Engineering: Creating the right features from raw data is crucial for predictive models to perform well.**

- **Model Evaluation: Use metrics like accuracy, precision, recall, F1 score (for classification), or RMSE (Root Mean Squared Error) and MAE (Mean Absolute Error) for regression tasks to evaluate how well the model performs.**

- **Overfitting: Make sure the model generalizes well and doesn't just memorize the training data (leading to poor performance on unseen data).**

Decision Making

Decision making uses gathered facts, analysis, and judgment. Decision making identifies an optimal solution for a problem or situation.

Once predictions are made, decision-making comes into play. In a data science context, decision-making is the process of selecting the best course of action based on the predictions and insights derived from data. It's about using predictive models to guide strategic or operational decisions.

Methods for Decision Making:

1. **Optimization: Once you have predictive models, you can use them in optimization problems. This involves making decisions that maximize or minimize some objective, given constraints.**

    - **Linear Programming: Solving optimization problems in operations research (e.g., maximizing profit, minimizing cost).**
    - **Integer Programming: Useful in scheduling or resource allocation where decisions are binary (yes/no).**

2. **A/B Testing: A/B testing is a way to make decisions based on experimental data. You create two (or more) variations of a product or process and see which performs better. This is a form of empirical decision-making, where the decision is informed by real-world data.**

    - **Null Hypothesis Testing: Statistical tests (e.g., t-tests, chi-squared) help assess whether differences between groups are statistically significant.**

3. **Recommendation Systems: Based on user behavior (such as product interactions or ratings), recommendation algorithms make decisions about what products or services to recommend to users. Common algorithms include:**
    - **Collaborative Filtering (based on user-item interactions).**
    - **Content-Based Filtering (based on product features).**
    - **Matrix Factorization (e.g., SVD, ALS).**

4. **Multi-Criteria Decision Analysis (MCDA): When there are multiple competing objectives or criteria (e.g., cost, quality, time), MCDA helps in making decisions that balance these criteria. It involves techniques like Weighted Scoring Models or Analytic Hierarchy Process (AHP).**

5. **Reinforcement Learning: In dynamic decision environments (like robotics or game AI), reinforcement learning helps make decisions by interacting with an environment and learning from feedback (rewards/punishments). This is useful for problems where you need to optimize sequential decisions over time, like inventory management or real-time bidding.**

6. **Game Theory: When multiple parties are involved, each making decisions based on what others are doing, game theory models can help predict optimal strategies and outcomes. For example, it can be used in competitive marketing, pricing, and auctions.**