




TARUN KAUSHIK





TARUN KAUSHIK

# SYLLABUS

5/8  
Que

## PAPER-I

250  
Marks

## PAPER-II



5/8  
Que

### 1. **LINEAR ALGEBRA**

2. Calculus
3. Ordinary Differential Equation
4. Analytical Geometry
5. Vector Analysis
6. Dynamics & Statics

1. Abstract Algebra
2. Real Analysis
3. **Complex Analysis**
4. **Linear Programming**
5. Partial Differential Equations
6. Numerical Analysis
7. Mechanics & Fluid Dynamics

I.



## Contents



TARUN KAUSHIK

*Preface*

*Syllabus (MDU, Rohtak and KU, Kurukshetra)*

*Basic Concepts*

*Chapters*

*Pages*

### SECTION-A Infinite Series

1. Infinite Series	...	3
--------------------	-----	---

### SECTION-B Matrices and Its Applications

*2. Matrices and Its Applications (Not for KUK)	...	73
---	-----	----

### SECTION-C Differential Calculus

*3. Successive Differentiation (Not for KUK)	...	131
*4. Leibnitz's Theorem and Applications (Not for KUK)	...	151
5. Taylor's and Maclaurin's Series	...	165
6. Curvature	...	183
7. Asymptotes	...	214
8. Curve Tracing	...	234
9. Functions of Two or More Variables (Limit and Continuity)	...	253
10. Partial Differentiation	...	262

### SECTION-D Integral Calculus

11. Applications of Single Integration (Volumes and Surfaces of Solids of Revolution)	...	341
12. Multiple Integrals	...	362


### SECTION-E Vector Calculus

**13. Vector Calculus (Not for MDU)	...	413
Appendix-I: Short Answer Type Questions	...	497
Appendix-II: Some Important Curves	...	510
Examination Papers	...	513

\*Chapters 2, 3 and 4 are not included in the syllabus of KU, Kurukshetra.

\*\*Chapter 13 is not included in the syllabus of MDU, Rohtak.

2.



## CONTENTS



TARUN KAUSHIK

*Preface*

*Syllabus*

*Basic Concepts*

*Chapters*

*Pages*

### SECTION-A

*(Vector Calculus)*

1*. Vector Calculus ( <i>Not for K.U.K.</i> )	...	3
2**. Matrices and its Applications ( <i>Not for M.D.U.</i> )	...	85

### SECTION-B

*(Ordinary Differential Equations and its Applications)*

3. Exact Differential Equations	...	147
4. Applications of Differential Equations of First Order and First Degree	...	156
5. Linear Differential Equations	...	168
6. Applications of Linear Differential Equations	...	204

### SECTION-C

*(Laplace Transforms and its Applications)*

7. Laplace Transforms and its Applications	...	219
--	-----	-----

### SECTION-D

*(Partial Differential Equations and its Applications)*

8. Partial Differential Equations	...	273
9. Applications of Partial Differential Equations	...	324


Appendix-I: Short Answer Type Questions	...	352
---	-----	-----

Appendix-II: Examination Papers	...	362
---------------------------------	-----	-----

3.



TARUN KAUSHIK



CONTENTS	
Chapters	Pages
Syllabus	(iv)
1. Fourier Series	1
2. Fourier Transforms	45
3. Functions of a Complex Variable	102
4. Probability Distributions and Hypothesis Testing	194
5. Linear Programming	285
Examination Paper	379
Appendices	383

# Linear Algebra Syllabus

- Vector spaces over  $\mathbb{R}$  and  $\mathbb{C}$ , linear dependence and independence, subspaces, bases, dimension; Linear transformations, rank and nullity,.
- Algebra of Matrices; Row and column reduction, Echelon form, congruence's and similarity; Rank of a matrix; Inverse of a matrix;
- Solution of system of linear equations; Eigen values and eigenvectors, characteristic polynomial, Cayley- Hamilton theorem,
- Symmetric, skew symmetric, Hermitian, skew-Hermitian, orthogonal and unitary matrices and their Eigen values.





TARUN KAUSHIK

## Matrix

### Linear Algebra

Dated: 10/03/2016

#### # Linearly Independent :-

Que: (2015-Mains) The vector: (10 Marks)

$V_1 = (1, 1, 2, 4)$ ;  $V_2 = (2, -1, -5, 2)$ ,  $V_3 = (1, -1, 4, 0)$  and  $V_4 = (2, 1, 1, 6)$  are linearly independent/dependent.

Sol.

$$\begin{bmatrix} 1 & 1 & 2 & 4 \\ 2 & -1 & -5 & 2 \\ 1 & -1 & -4 & 0 \\ 2 & 1 & 1 & 6 \end{bmatrix} \xrightarrow{\substack{R_2 \leftrightarrow R_2 - 2R_1 \\ R_3 \leftrightarrow R_3 - R_1 \\ R_4 \leftrightarrow R_4 - 2R_1}} \begin{bmatrix} 1 & 1 & 2 & 4 \\ 0 & -3 & -9 & -6 \\ 0 & -2 & -6 & -4 \\ 0 & -1 & -3 & -2 \end{bmatrix} \xrightarrow{\substack{R_2 \leftrightarrow R_2 \\ R_3 \leftrightarrow R_3 \\ R_4 \leftrightarrow R_4 - R_2}} \begin{bmatrix} 1 & 1 & 2 & 4 \\ 0 & -3 & -9 & -6 \\ 0 & -2 & -6 & -4 \\ 0 & -1 & -3 & -2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 2 & 4 \\ 0 & 1 & 3 & 2 \\ 0 & 1 & 3 & 2 \\ 0 & -1 & -3 & -2 \end{bmatrix} \xrightarrow{\substack{R_2 \leftrightarrow R_2 - R_1 \\ R_4 \leftrightarrow R_4 + R_2}} \begin{bmatrix} 1 & 1 & 2 & 4 \\ 0 & 1 & 3 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

⇒ Hence this shows that the vector  $V_1$  and  $V_2$  are linearly independent vectors &  $V_3$  &  $V_4$  are linearly dependent.

(Mains 2015)

Que:- Reduce the following matrix to the Row Echelon form and hence find its Rank.

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 2 & 1 & 4 & 5 \\ 1 & 5 & 5 & 7 \\ 8 & 1 & 14 & 17 \end{bmatrix}$$

Sol.  $R_2 \leftrightarrow R_2 - 2R_1$ ,  $R_3 \leftrightarrow R_3 - R_1$ ,  $R_4 \leftrightarrow R_4 - 8R_1$

$$= \begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & -3 & -2 & -3 \\ 0 & 3 & 2 & 3 \\ 0 & -15 & -10 & -15 \end{bmatrix} \quad \text{Now } R_4 \leftrightarrow R_4 - 5R_2$$

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & -3 & -2 & -3 \\ 0 & 3 & 2 & 3 \\ 0 & 3 & 2 & 3 \end{bmatrix} \xrightarrow{\substack{R_3 \leftrightarrow R_3 - R_2 \\ R_4 \leftrightarrow R_4 - R_2}} \begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & -3 & -2 & -3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Rank = 2

↑ Reduced Echelon matrix.

Que:- (Mains 2015)

(12 Marks)

If  $A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$ ; find  $A^{30}$ .

Sol.

Using Cayley Hamilton theorem.

$$|A - \lambda I| = 0$$

$$\begin{vmatrix} 1-\lambda & 0 & 0 \\ 1 & -\lambda & 1 \\ 0 & 1 & -\lambda \end{vmatrix} = (1-\lambda)(\lambda^2-1) = 0$$

$$3 - \lambda^2 - \lambda + 1 = 0$$

$$A^3 - A^2 - A + I = 0$$

$$A^3 = A^2 + A - I$$

$$A^4 = A^3 + A^2 - A = A^2 + A - I + A^2 - A = 2A^2 - I$$

$$A^8 = 4A^4 + I - 4A^2 = 4(2A^2 - I) + I - 4A^2 = 4A^2 - 3I$$

$$A^{16} = 16A^8 + 9I - 24A^2 = 32A^2 - 16I + 9I - 24A^2 = 8A^2 - 7I$$

$$A^{32} = 64A^{16} + 49I - 112A^2 = 128A^2 - 64I + 49I - 112A^2$$

$$\begin{aligned} A^4 &= 2A^2 - I & A^{64} &= 32A^2 - 31I \\ A^8 &= 4A^2 - 3I & A^{128} &= 64A^2 - 63A \\ A^{16} &= 8A^2 - 7I & A^{256} &= 128A^2 - 127I \\ A^{32} &= 16A^2 - 15I & A^{512} &= 256A^2 - 255I \end{aligned}$$

$$= 16A^2 - 15I$$

$$\Rightarrow A^{30} = 16A^2 - 15A^{-2}$$





TARUN KAUSHIK

$$A^3 = A + A - I \Rightarrow A = I + A^1 - A^2$$

$$\Rightarrow A^2 = I + A^1 - A$$

$$\rightarrow A^3 = 16I - 15A^2$$

$$= 16I - 15(I + A^1 - A) = 16I - 15I + 15(A - A^1)$$

$$A^3 = I + 15(A - A^1)$$

Shortcut to Calculate Inverse of a matrix:-

$$\begin{array}{c|cccc} 1 & 0 & 0 & 1 & 0 \\ \hline 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 1 & 0 \end{array}$$

$$\begin{bmatrix} -1 & 1 & 0 \\ 0 & 0 & -1 \\ 1 & -1 & 0 \end{bmatrix} = \text{Adj}(A)$$

$$A^{-1} = \frac{1}{|A|} \begin{bmatrix} -1 & 1 & 0 \\ 0 & 0 & -1 \\ 1 & -1 & 0 \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 0 & 1 \\ -1 & 1 & 0 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$|A| = -1$$

$$A^{-1} = 15 \left( \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} - \begin{bmatrix} 1 & -1 & 0 \\ 0 & 0 & 1 \\ -1 & 1 & 0 \end{bmatrix} \right) + I$$

$$= 15 \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 15 & 0 \\ 15 & 0 & 0 \\ 15 & 0 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 15 & 0 \\ 15 & 1 & 0 \\ 15 & 0 & 1 \end{bmatrix} \text{ Ans}$$

Que (Mains 2015) (12 Marks)

$\{(1,0,0,0), (0,1,0,0), (1,2,0,1), (0,0,0,1)\}$  ← vector

i) basis

ii) dimension

Ans

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 2 & 0 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$R_3 \leftrightarrow R_3 - R_1$$

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 2 & 0 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\rightarrow R_3 \leftrightarrow R_3 - 2R_2$$

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$R_4 \leftrightarrow R_4 - R_3$$

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\text{Basis} = \{(1,0,0,0), (0,1,0,0), (0,0,0,1)\}$$

$$\dim = 3. \text{ Ans}$$

Que (Mains 2014) (10 Marks)

System of Linear Equation:-

$$x + y + z = 6$$

$$x + 2y + 3z = 10$$

$$x + 2y + 4z = 11$$

i) No solution

ii) unique solution

iii) infinite No. of sol.

Sol

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \\ 1 & 2 & 4 \end{bmatrix}$$

$$B = \begin{bmatrix} 6 \\ 10 \\ 11 \end{bmatrix}$$

$$[A:B]$$

$$[A:B] = \begin{bmatrix} 1 & 1 & 1 & : & 6 \\ 1 & 2 & 3 & : & 10 \\ 1 & 2 & 4 & : & 11 \end{bmatrix}$$

$$R_2 \leftrightarrow R_2 - R_1$$

$$R_3 \leftrightarrow R_3 - R_1$$

$$= \begin{bmatrix} 1 & 1 & 1 & : & 6 \\ 0 & 1 & 2 & : & 4 \\ 0 & 1 & 3 & : & 5 \end{bmatrix}$$

$$R_3 \leftrightarrow R_3 - R_2$$

$$= \begin{bmatrix} 1 & 1 & 1 & : & 6 \\ 0 & 1 & 2 & : & 4 \\ 0 & 0 & 1 & : & 1 \end{bmatrix}$$





TARUN KAUSHIK

Case: I (No solution)  
 $\rho(A) \neq \rho(A:B)$

for this:

$$\lambda = 3, \mu \neq 10$$

ii) unique solution:  $\mu$

$$\lambda \neq 3, \mu \neq 10$$

(iii) Many solution:-

$$\lambda = 3 \text{ and } \mu = 10$$

①  $\rho(A) = \rho(A:B) = \text{No. of variable}$   
 $\Rightarrow$  unique solution

②  $\rho(A) \neq \rho(A:B)$   
 $\Rightarrow$  No sol.

③  $\rho(A) = \rho(A:B) < \text{No. of variable}$   
 $\Rightarrow$  Many solution

Que: (2014) (8-Marks) "Eigen Values & Eigen Vector"

$$A = \begin{bmatrix} -2 & 2 & -3 \\ 2 & 1 & -6 \\ -1 & -2 & 0 \end{bmatrix}$$

Sol.

$$|A - \lambda I| = 0; \quad \begin{vmatrix} -2-\lambda & 2 & -3 \\ 2 & 1-\lambda & -6 \\ -1 & -2 & -\lambda \end{vmatrix}$$

$$A = \begin{bmatrix} -2 & 2 & -3 \\ 2 & 1 & -6 \\ -1 & -2 & 0 \end{bmatrix}$$

$$|A - \lambda I| = 0$$

$$\begin{bmatrix} -2-\lambda & 2 & -3 \\ 2 & 1-\lambda & -6 \\ -1 & -2 & -\lambda \end{bmatrix}$$

$$44 \times \frac{5}{2}$$

$$\begin{aligned} & (-2-\lambda) [(1-\lambda)(-\lambda) + 12] - 2[-2\lambda - 6] - 3[-4 + 1 - \lambda] \\ & + (2+\lambda)(1-\lambda^2 + 12) + 4\lambda + 12 + 9 + 3\lambda \\ & 2\lambda - 2\lambda^2 + 24 + \lambda^2 - \lambda^3 + 12\lambda \\ & = -\lambda^3 - \lambda^2 + 19\lambda + 45 = 0 \end{aligned}$$





TARUN KAUSHIK

$$\lambda^3 + \lambda^2 - 21\lambda - 45 = 0 \quad (\lambda = -3)$$

$$(\lambda + 3)(\lambda^2 - 2\lambda - 15) = 0 \quad \begin{pmatrix} -2 & 2 & -3 \\ 2 & 1 & -6 \\ -1 & -2 & 0 \end{pmatrix}$$

$$\lambda = -3, -3, 5$$

for  $\lambda = -3$

$$\begin{pmatrix} 1 & 2 & -3 \\ 2 & 4 & -6 \\ -1 & -2 & 3 \end{pmatrix}$$

$$\begin{bmatrix} 1 & 2 & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$\begin{pmatrix} -7 & 2 & -3 \\ 2 & -4 & -6 \\ -1 & -2 & -5 \end{pmatrix}$$

$$\begin{bmatrix} +1 & +2 & +5 \\ 2 & -4 & -6 \\ -7 & 2 & -3 \end{bmatrix}$$

$$\begin{pmatrix} 1 & 2 & 5 \\ 0 & -8 & -16 \\ 0 & +16 & 32 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 2 & 5 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix}$$

$$x_1 + 2x_2 + 5x_3 = 0; \quad x_2 + 2x_3 = 0$$

$$x_2 = -2x_3$$

JAN-July

$$x_1 - 4x_3 + 5x_3 = 0$$

$$x_1 = x_3$$

$$\frac{x_1}{1} = \frac{x_2}{-2} = \frac{x_3}{1}$$

★ Orthogonal Matrix :

$$\text{if } A \cdot A^T = I \Rightarrow \uparrow$$

★ Hermitian Matrix :



$$a_{ij} = \bar{a}_{ji}$$

Complex No. Conjugate

Skew-Hermitian Matrix :-

$$a_{ij} = -\bar{a}_{ji}$$

Complex Numbers  
Conjugate





TARUN KAUSHIK

(12 marks)  
Que:- (2015) Find Eigen values & eigen vectors of Matrix

$$\begin{bmatrix} 1 & 1 & 3 \\ 1 & 5 & 1 \\ 3 & 1 & 1 \end{bmatrix} = A$$

Solution  $[A - \lambda I] = 0 \Rightarrow \begin{bmatrix} 1-\lambda & 1 & 3 \\ 1 & 5-\lambda & 1 \\ 3 & 1 & 1-\lambda \end{bmatrix} = [A - \lambda I]$

$$|A - \lambda I| = 0$$

$$\begin{aligned} &\Rightarrow (1-\lambda)[(5-\lambda)(1-\lambda)-1] - 1[(1-\lambda)-3] + 3(1-15+3\lambda) \\ &= (1-\lambda)(5-5\lambda-\lambda+\lambda^2-1) - (-2-\lambda) + 3(-14+3\lambda) \\ &= (1-\lambda)(4-6\lambda+\lambda^2) + 2+\lambda-42+9\lambda \\ &= 4-6\lambda+\lambda^2-4\lambda+6\lambda^2-\lambda^3+2+\lambda-42+9\lambda \\ &= -\lambda^3+7\lambda^2-36=0 \Rightarrow \lambda^3-7\lambda^2+36=0 \end{aligned}$$

by trial method  $\lambda = -2$  is a factor.

$$\begin{array}{r} (\lambda+2) \overline{\lambda^3-7\lambda^2+36} \quad \lambda^2-9\lambda+18 \\ \underline{\lambda^3+2\lambda^2} \phantom{+36} \\ -9\lambda^2+36 \phantom{+18\lambda} \\ \underline{-9\lambda^2+18\lambda} \phantom{+36} \\ 18\lambda+36 \\ \underline{18\lambda+36} \\ 0 \end{array}$$

$$\Rightarrow \text{Eigen value} = \lambda = -2, +3, +6$$

Eigen vectors:-

i) for  $\lambda = -2 \Rightarrow [A - \lambda I] = \begin{bmatrix} 3 & 1 & 3 \\ 1 & 7 & 1 \\ 3 & 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$

$$R_3 \leftrightarrow R_3 - R_1 \Rightarrow \begin{bmatrix} 3 & 1 & 3 \\ 1 & 7 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0$$

$$R_2 \leftrightarrow R_2 - 3R_1$$

$$\begin{bmatrix} 3 & 1 & 3 \\ 0 & 20 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0$$

$$\Rightarrow 3x_1 + x_2 + 3x_3 = 0 \quad \text{and} \quad x_2 = 0$$

$$3x_1 + 3x_3 = 0 \Rightarrow 3x_1 = -3x_3 \\ x_1 = -x_3$$

vector  $(1, 0, -1)$

ii) for  $\lambda = 3$

$$[A - \lambda I] = \begin{bmatrix} -2 & 1 & 3 \\ 1 & 2 & 1 \\ 3 & 1 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$R_1 \leftrightarrow R_2$$

$$\begin{bmatrix} 1 & 2 & 1 \\ -2 & 1 & 3 \\ 3 & 1 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0$$

$$R_2 \leftrightarrow R_2 + 2R_1, R_3 \leftrightarrow R_3 - 3R_1$$

$$\begin{bmatrix} 1 & 2 & 1 \\ 0 & 5 & 5 \\ 0 & -5 & -5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0$$

$$R_3 \leftrightarrow R_3 + R_2$$

$$\begin{bmatrix} 1 & 2 & 1 \\ 0 & 5 & 5 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0$$

$$x_1 + 2x_2 + x_3 = 0$$

$$x_2 + x_3 = 0$$

$$x_1 = x_3$$

$$x_2 = -x_3$$

vector  $(1, -1, 1)$

iii) for  $\lambda = 6$

$$\begin{bmatrix} -5 & 1 & 3 \\ 1 & -1 & 1 \\ 3 & 1 & -5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0$$

$$R_1 \leftrightarrow R_2$$

$$\begin{bmatrix} 1 & -1 & 1 \\ -5 & 1 & 3 \\ 3 & 1 & -5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0$$

$$R_2 \leftrightarrow R_2 + 5R_1, R_3 \leftrightarrow R_3 - 3R_1$$

$$\begin{bmatrix} 1 & -1 & 1 \\ 0 & -4 & 8 \\ 0 & 4 & -8 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0$$

$$R_3 \leftrightarrow R_3 + R_2$$

$$\begin{bmatrix} 1 & -1 & 1 \\ 0 & -4 & 8 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0$$

$$x_1 - x_2 + x_3 = 0; -4x_2 + 8x_3 = 0$$

$$x_1 = x_3$$

$$x_2 = 2x_3$$

vector  $(1, 2, 1)$