



TARUN KAUSHIK

# MATHEMATICS FOR UPSC CSE MAINS (LAPLACE TRANSFORM)

Group "Maths as Optional for IAS" +91-9416421592





TARUN KAUSHIK

## Laplace Transform

Dated: 30/08/2016

①

→ L.T: Mathematical tool to convert a time domain signal into frequency domain signal.

Reason:- Since; it is quite easy to analyse the signal in time domain but still why we transform it into frequency domain. Because it is easy to do the calculation in frequency domain than time domain.

# Fourier Series | Fourier Transform | L.T

Periodic

Aperiodic

$$L[x(t)] = \int_{-\infty}^{\infty} x(t) e^{-st} dt$$

t: time

s: Complex frequency

jω (imaginary)

σ (Real)

$$s = \sigma + j\omega$$

when  $\sigma = 0$

$$s = j\omega$$

Complex freq.

$$F[x(t)] = \int_{-\infty}^{\infty} x(t) e^{-j\omega t} dt$$

So F.T is a special case of L.T. when the Real part of Complex freq. is zero.

$$L[x(t)] = \int_{-\infty}^{\infty} x(t) \cdot e^{-st} dt$$

Some Imp. formulas:-

②

$$① \quad L[1] = \frac{1}{s}$$

$$② \quad L[t] = n! / s^{n+1}$$

$$③ \quad L[\sin at] = \frac{a}{s^2 + a^2}$$

$$④ \quad L[\cos at] = \frac{s}{s^2 + a^2}$$

$$⑤ \quad L[e^{at}] = \frac{1}{s-a}$$

$$⑥ \quad L[\sin hat] = \frac{a}{s^2 + a^2}$$

$$⑦ \quad L[\cos hat] = \frac{s}{s^2 + a^2}$$

# Properties:-

① Linearity Property:-

$$L[ax_1(t) + bx_2(t)] = aL[x_1(t)] + bL[x_2(t)]$$

→ "Homogeneity & super-position"

② Shifting Property:-

$$\text{Time: } L[x(t-t_0)] = e^{-st_0} \cdot X(s)$$

$$\text{Freq: } L[e^{s_0 t} x(t)] = X(s-s_0)$$

$$L[e^{at} \sin bt]$$

$$\frac{b}{s^2 + b^2}$$

$$\frac{b}{(s-a)^2 + b^2} \text{ Ans.}$$

③ Time Scaling:-

$$x(t) \longleftrightarrow X(s)$$

$$x(at) \longleftrightarrow \frac{1}{|a|} X\left(\frac{s}{a}\right)$$

(Exp)

(Comp)



Que: - (1)  $L[t \cos t]$  ; (2)  $L[t \sin t]$

(3)  $L[t^2 \sin at]$

Que: -  $L[t e^{-4t} \sin 3t]$

Sol.<sup>n</sup>  $L[t] = \frac{1}{s^2}$   $L[t e^{+i3t}] = \frac{1}{(s-i3)^2} \times \frac{(s+i3)^2}{(s-i3)^2}$

$$\left[ \because e^{it} = \cos t + i \sin t \right]$$
$$L[t e^{i3t}] = \frac{(s^2 - 9) + i6s}{(s^2 + 9)^2}$$
$$L[t (\cos 3t + i \sin 3t)] = "$$

(4)  $\Rightarrow L[t \sin 3t] = \frac{6s}{(s^2 + 9)^2}$

Now  $L[e^{-4t} \cdot t \sin 3t]$

$$\begin{aligned} &\xrightarrow{\quad} \frac{6s}{(s^2 + 9)^2} \\ &\xrightarrow{\quad} \frac{6(s+4)}{((s+4)^2 + 9)^2} \text{ Ans.} \end{aligned}$$

Que:  $f(t) = |t-1| + |t+1| + |t+2| + |t-2|$ ;  $t \geq 0$

Sol.<sup>n</sup>

$$\begin{aligned} t-1=0 &\Rightarrow t=1 \\ t+1=0 &\Rightarrow t=-1 \\ t+2=0 &\Rightarrow t=-2 \\ t-2=0 &\Rightarrow t=2 \end{aligned} \quad \times \because t \geq 0$$

so interval will be  $0 \leq t \leq 1$  :-  $f_1(t)$

$1 \leq t \leq 2$  :  $f_2(t)$

$t > 2$  :  $f_3(t)$

$$f_1(t) = -(t-1) + t+1 + t+2 - (t-2) = 6$$

$$f_2(t) = t-1 + t+1 + t+2 - (t-2) = 2t+4$$

$$f_3(t) = (t-1) + (t+1) + (t+2) + (t-2) = 4t$$

$$\therefore L[f(t)] = \int_0^{\infty} f(t) e^{-st} dt$$

$$= \int_0^1 6 \cdot e^{-st} dt + \int_1^2 (2t+4) e^{-st} dt + \int_2^{\infty} 4t \cdot e^{-st} dt$$

$$= \frac{1}{s^2} [6s + 2e^{-s} + 2e^{-2s}]$$

$$= \frac{2}{s^2} (3s + e^{-s} + e^{-2s}) \text{ Ans.}$$





⑥

Que:- I.L.T.  $X(s) = \log \frac{s+1}{s-1}$ ;

Sol.<sup>n</sup>

$$X(s) = L[x(t)] = \log \frac{(s+1)}{(s-1)}$$

$$L[t \cdot x(t)] = (-1) \frac{d}{ds} X(s)$$

$$= (-1) \frac{d}{ds} \left( \log \frac{s+1}{s-1} \right)$$

$$= (-1) \frac{d}{ds} [\log(s+1) - \log(s-1)]$$

$$= - \left[ \frac{1}{s+1} - \frac{1}{s-1} \right] = - \left[ \frac{s-1-s-1}{s^2-1} \right]$$

$$L[t \cdot x(t)] = \frac{2}{s^2-1} = 2 \left( \frac{1}{s^2-1} \right)$$

$$= 2 L[\sinh t]$$

$$t \cdot x(t) = 2 \sinh t$$

$$\Rightarrow x(t) = \frac{2 \sinh t}{t} \text{ Ans.}$$

Que:- L.T. of  $\frac{e^{-at} - e^{-bt}}{t}$

Sol.<sup>n</sup>

Similar like integral property (freq.<sup>n</sup>)

$$L \left[ \frac{e^{-at} - e^{-bt}}{t} \right] = \left[ \frac{1}{s+a} - \frac{1}{s+b} \right]$$

$$L \left( \frac{x(t)}{t} \right) \iff \int_s^\infty X(s) ds$$

$$= \int_s^\infty \frac{1}{s+a} - \frac{1}{s+b} ds = \log \frac{s+a}{s+b} \Big|_s^\infty$$

$$= \log \frac{s(1+a/s)}{s(1+b/s)} \Big|_s^\infty$$

$$= \log 1 - \log \frac{s+a}{s+b} = \boxed{\log \frac{s+b}{s+a}} \text{ Ans.}$$

⑤

④ Differentiation :-

$$\begin{array}{l} \text{Time : } x(t) \longleftrightarrow X(s) \\ \frac{d}{dt} x(t) \longleftrightarrow s \cdot X(s) - x(0) \end{array}$$

$$\begin{array}{l} \text{Freq. : } x(t) \longleftrightarrow X(s) \\ -t \cdot x(t) \longleftrightarrow \frac{d}{ds} X(s) \end{array}$$

⑤ Integral :-

$$\begin{array}{l} \text{Time :- } x(t) \longleftrightarrow X(s) \\ \int_0^t x(\tau) d\tau \longleftrightarrow \frac{X(s)}{s} \end{array}$$

$$\begin{array}{l} \text{Freq. :- } x(t) \longleftrightarrow X(s) \\ \frac{x(t)}{t} \longleftrightarrow \int_s^\infty X(s) ds \end{array}$$

Using different

$$\begin{cases} L[x(t)] = X(s) \\ L[x'(t)] = sX(s) - x(0) \\ L[x''(t)] = s^2X(s) - sx(0) - x'(0) \\ L[x'''(t)] = s^3X(s) - s^2x(0) - sx'(0) - x''(0) \\ \vdots \\ L[x^{(n)}(t)] = s^nX(s) - s^{n-1}x(0) - s^{n-2}x'(0) - \dots \end{cases}$$

⑥ multiplication by "t<sup>n</sup>" :-

$$L[t^n \cdot x(t)] = (-1)^n \frac{d^n}{ds^n} X(s) \quad (n \in \mathbb{N})$$

Eg.  $L[t \cdot \sin^2 3t] = ?$

$$\sin^2 3t = \frac{1 - \cos 6t}{2}$$

$$L \left[ \frac{1 - \cos 6t}{2} \right] = \frac{1}{2} \left( \frac{1}{s} - \frac{1}{s^2 + 36} \right)$$

$$(-1) \frac{d}{ds} \left( \frac{1}{s(s^2+36)} \right) = \frac{54(s^2+12)}{s^2(s^2+36)^2} \text{ Ans.}$$



(7)

Convolution Property :-  $x(t) \leftrightarrow X(s)$   
 $h(t) \leftrightarrow H(s)$

$$x(t) * h(t) \leftrightarrow X(s) \cdot H(s)$$

Proof.  $L[x(t) * h(t)] = \int_{-\infty}^{\infty} [x(t) * h(t)] e^{-st} dt$

[But  $x(t) * h(t) = \int_0^t x(\tau) h(t-\tau) d\tau$ ]


$$= \int_0^{\infty} \int_0^t x(\tau) h(t-\tau) e^{-st} d\tau dt$$

on changing the Order of Integration

means:-

Original  $(0 < \tau < t)$  LR

After changing  $(0 < \tau < \infty)$  BT



L-R	T
	1
	B

$$= \int_0^{\infty} \int_{\tau}^{\infty} x(\tau) h(t-\tau) e^{-st} dt d\tau$$

Put  $t-\tau = v$   
 $t = v + \tau$

$$= \int_0^{\infty} \int_0^{\infty} x(\tau) h(v) e^{-s(v+\tau)} dv d\tau$$

$$= \int_0^{\infty} x(\tau) e^{-s\tau} d\tau \cdot \int_0^{\infty} h(v) e^{-sv} dv$$

$$= X(s) \cdot H(s) \text{ Proved.}$$

# Inverse Laplace Transform :-

(8)

(1)  $L^{-1}\left(\frac{1}{s^n}\right) = \frac{t^{n-1}}{(n-1)!}$

(2)  $L^{-1}\left(\frac{1}{(s-a)^n}\right) = \frac{t^{n-1}}{(n-1)!} e^{at}$

Que:-  $L^{-1}\left\{\frac{s^2}{(s^2+a^2)(s^2+b^2)}\right\}$

Soln  $\frac{s^2}{(s^2+a^2)(s^2+b^2)} = \frac{1}{(s^2+a^2)} \times \frac{s}{(s^2+b^2)} = X(s) \cdot H(s)$

Conv. Prop. :  $x(t) * h(t) = L^{-1}[X(s) \cdot H(s)]$

$X(s) = \frac{1}{s^2+a^2} \Rightarrow x(t) = \cos at$

$H(s) = \frac{1}{s^2+b^2} \Rightarrow h(t) = \cos bt$

$x(t) * h(t) = \int_0^t x(\tau) h(t-\tau) d\tau$

Ans =  $\frac{1}{2} \int_0^t 2 \cos a\tau \cos b(t-\tau) d\tau$   
 $= \frac{1}{2} \int_0^t [\cos(a\tau + b(t-\tau)) + \cos(a\tau - b(t-\tau))] d\tau$   
 $= \frac{1}{2} \int_0^t [\cos((a-b)\tau + bt) + \cos((a+b)\tau - b\tau)] d\tau$   
 $= \frac{1}{2} \left[ \frac{\sin((a-b)\tau + bt)}{a-b} + \frac{\sin((a+b)\tau - b\tau)}{a+b} \right]_0^t$   
 $= \frac{1}{2} \left( \frac{\sin at - \sin bt}{a-b} + \frac{\sin at + \sin bt}{a+b} \right)$  Ans





TARUN KAUSHIK

Q.9.

Que. - Evaluate integral :-  $\int_0^{\infty} \left[ \frac{e^{-t} - e^{-3t}}{t} \right] dt$

$$= \int_0^{\infty} e^{-ot} \left( \frac{e^{-t} - e^{-3t}}{t} \right) dt \Rightarrow \text{Indirectly means to find the L.T. of}$$

Let  $x(t) = e^{-t} - e^{-3t}$   
 $\Rightarrow X(s) = \left( \frac{1}{s+1} - \frac{1}{s+3} \right)$

Prop. :  $\frac{x(t)}{t} \xrightarrow{LT} \int_1^{\infty} X(s) ds$

$$L\left[\frac{x(t)}{t}\right] = \int_1^{\infty} \left( \frac{1}{s+1} - \frac{1}{s+3} \right) ds$$

$$= \log(s+1) - \log(s+3) \Big|_1^{\infty}$$

$$= \log \frac{s+1}{s+3} \Big|_1^{\infty} = \log 1 - \log \frac{s+1}{s+3}$$

$$= \boxed{\log \frac{s+3}{s+1}} \quad \text{Here } s=0$$

$$\Rightarrow \int_0^{\infty} \frac{e^{-t} - e^{-3t}}{t} dt = \boxed{\log 3} \text{ Ans.}$$

Q(2015) : 6 Marks :- I.L.T.

$$L^{-1} \left\{ \log \left( 1 + \frac{1}{s^2} \right) + \frac{s}{s^2+25} e^{-\pi s} \right\}$$

$$\stackrel{\text{Sol.}^n}{=} L^{-1} \left\{ \log \left( 1 + \frac{1}{s^2} \right) \right\} + L^{-1} \left\{ \frac{s}{s^2+25} e^{-\pi s} \right\}$$

$$\text{I : } L[t \cdot x(t)] = - \frac{d}{ds} (\log(s^2+1) - \log s^2)$$

$$= - \left( \frac{2s}{s^2+1} - \frac{1}{s^2} \right) = \frac{1}{s^2} - \frac{2s}{s^2+1}$$

$$\Rightarrow \boxed{x(t) = 1 - 2 \cos t}$$

$$\text{II : } L^{-1} \left\{ \frac{s}{s^2+25} e^{-\pi s} \right\}$$

Shift<sup>n</sup> Prop. :  $x(t) \longleftrightarrow X(s)$   
 $x(t-t_0) \longleftrightarrow e^{-st_0} X(s)$

Here  $X(s) = \frac{s}{s^2+25} \Rightarrow x(t) = \cos 5t$

But using shifting property :-

$$\Rightarrow \cos 5(t-\pi) = \cos(5\pi-5t) = -\sin 5t$$

$$\text{so } L^{-1} \left\{ \log \left( 1 + \frac{1}{s^2} \right) + \frac{s e^{-\pi s}}{s^2+25} \right\} = \left[ 1 - 2 \cos t - \sin 5t \right] \text{ Ans.}$$

Q(2015) : 6 marks :  $y'' + y = t$

Sol.<sup>n</sup> taking L.T. both sides  $y(0)=1$   
 $y'(0)=-2$

$$[s^2 Y(s) - s y(0) - y'(0)] + Y(s) = \frac{1}{s^2}$$

$$\Rightarrow s^2 Y(s) - s + 2 + Y(s) = \frac{1}{s^2}$$

$$\Rightarrow Y(s) (1+s^2) = \frac{1}{s^2} + s - 2$$

$$\Rightarrow Y(s) = \frac{1}{s^2(s^2+1)} + \frac{s}{s^2+1} - \frac{2}{s^2+1}$$

$$\Rightarrow y(t) = [t - \sin t] + \cos t - 2 \sin t$$

$$= \boxed{t + \cos t - 3 \sin t} \text{ Ans.}$$

\* Prop.

$$t x(t) \longleftrightarrow X(s)$$

$$\int_0^t x(t) \longleftrightarrow \int_0^s X(s) ds$$

$$\int_0^t 1 - \cos t dt = t - \sin t \Big|_0^t = t - \sin t$$

$$x(t) = \sin t$$

$$\int_0^t \sin t dt = -\cos t \Big|_0^t = -\cos t + 1$$

(11)

Q (2014): 20 Marks: Initial Value Problem

$$\frac{d^2 y}{dt^2} + y = 8e^{-2t} \sin t; \quad y(0) = y'(0) = 0$$

Sol.<sup>n</sup>  $y'' + y = 8e^{-2t} \sin t$

Taking L.T. on both sides

$$\Rightarrow [s^2 Y(s) - s y(0) - y'(0)] + Y(s) = \frac{8}{(s+2)^2 + 1}$$

$$\Rightarrow (s^2 + 1) Y(s) = \frac{8}{(s+2)^2 + 1} = \frac{8}{s^2 + 4s + 5}$$

$$\Rightarrow Y(s) = \frac{8}{(s^2 + 4s + 5)(s^2 + 1)}$$

$$= \frac{As + B}{s^2 + 4s + 5} + \frac{Cs + D}{s^2 + 1}$$

$$\Rightarrow 8 = As^3 + Bs^2 + As + B + Cs^3 + 4Cs^2 + 5Cs + Ds^2 + 4Ds + 5D$$

Comparing coeff. of  $s^3, s^2, s^1, s^0$ ; we get

$$A + C = 0; \quad B + 4C + D = 0; \quad A + 5C + 4D = 0;$$

$$\Rightarrow A = -C; \quad B + 5D = 8$$

$$B - 3D = 0$$

$$-B + 5D = 8$$

$$-8D = -8 \Rightarrow D = 1$$

$$C = -1; A = 1$$

$$B = 3$$

$$\Rightarrow Y(s) = \frac{s+3}{s^2+4s+5} + \frac{-s+1}{s^2+1} = \frac{(s+2)+1}{s^2+4s+5} + \frac{-s+1}{s^2+1}$$

$$\Rightarrow Y(s) = \frac{(s+2)}{(s^2+2)^2+1} + \frac{1}{(s^2+2)^2+1} + \frac{3}{s^2+1} + \frac{1}{s^2+1}$$

$$\Rightarrow y(t) = e^{-2t} \cos t + e^{-2t} \sin t - \cos t + \sin t$$

(12)

Q: (2013); 15 Marks: Solve using L.T.

$$(D^2 + n^2)x = a \sin(nt + \alpha); \quad a, n, \alpha \text{ Constant}$$

$$x=0, \frac{dx}{dt}=0 \text{ at } t=0$$

Sol.<sup>n</sup> Taking L.T. on both sides  
 $s^2 X(s) - s x(0) - x'(0) + n^2 X(s) = L[a \sin(nt + \alpha)]$

$$\Rightarrow s^2 X(s) + n^2 X(s) = L[a (\sin nt \cos \alpha + \cos nt \sin \alpha)]$$

$$(s^2 + n^2) X(s) = \frac{an \cos \alpha}{s^2 + n^2} + \frac{s \sin \alpha}{s^2 + n^2}$$

$$\Rightarrow X(s) = \frac{an \cos \alpha}{(s^2 + n^2)^2} + \frac{\sin \alpha \cdot s}{(s^2 + n^2)^2}$$

Taking Inverse L.T.

$$\because L^{-1}\left[\frac{1}{(s^2 + a^2)^2}\right] = \frac{1}{2a^3} (\sin at - at \cos at)$$

$$\because L^{-1}\left\{\frac{s}{(s^2 + a^2)^2}\right\} = \frac{1}{2a} (t \sin at)$$

$$\Rightarrow x(t) = \frac{(an \cos \alpha)}{2n^3} (\sin nt - nt \cos nt) + \frac{1}{2n} (t \sin nt)$$

Ans

Q (2012): 12 Marks: Solve using L.T

$$y'' + 2y' + y = e^{-t}; \quad y(0) = -1, y'(0) = 1$$

Sol.<sup>n</sup>  $s^2 Y(s) - s y(0) - y'(0) + 2s Y(s) + 2Y(s) = \frac{1}{s+1}$

$$\Rightarrow s^2 Y(s) + s - 1 + 2s Y(s) + 2Y(s) = \frac{1}{s+1}$$



TARUN KAUSHIK



(13.)

$$\Rightarrow (s^2 + 2s + 1) Y(s) = \frac{1}{(s+1)} - (s+1)$$

$$\Rightarrow Y(s) = \frac{1}{(s+1)^3} - \frac{(s+1)}{(s+1)^3} = \frac{1}{(s+1)^3} - \frac{1}{(s+1)}$$

Taking Inverse L.T. :-

$$y(t) = e^{-t} \frac{t^2}{2} - e^{-t} = \boxed{e^{-t} \left( \frac{t^2}{2} - 1 \right)} \text{ Ans.}$$

Q: (2011) 15 Marks: Obtain General Sol.<sup>n</sup> of 2nd Order

O.D.E  $y'' + 2y' - 2y = x + e^x \cos x$  Take

Sol.<sup>n</sup> Taking L.T. on both sides; initial conditions zero.

$$s^2 Y(s) - 2sY(s) + 2Y(s) = \frac{1}{s} + \frac{(s-1)}{(s-1)^2 + 1}$$

$$(s^2 - 2s + 2) Y(s) = \frac{1}{s} + \frac{(s-1)}{(s-1)^2 + 1}$$

$$\Rightarrow Y(s) = \frac{1}{s((s-1)^2 + 1)} + \frac{(s-1)}{((s-1)^2 + 1)^2} \leftarrow \text{Convolut. Prop.}$$

Taking Inverse L.T. :-

$$\Rightarrow y(x) = \frac{e^x}{2} (\sin x - \cos x) + \frac{1}{2} + \frac{e^x}{2} \cdot x \sin x$$

$$\text{Prop.} \rightarrow \mathcal{L}^{-1} \left[ \frac{Y(s)}{s} \right] = \int_0^t x(t) dt$$

$$x(t) = \int_0^t \sin t dt$$

$$= \frac{e^t}{2} (\sin t - \cos t) \Big|_0^t$$

Q: (2011) 15 Marks Ans.

$$\frac{d^2 x}{dt^2} - 2 \frac{dx}{dt} + x = e^t$$

Sol.<sup>n</sup> Take L.T. on both sides  $x(0) = 2; \frac{dx}{dt} \Big|_{t=0} = 1$

$$\Rightarrow s^2 X(s) - sX(0) - x'(0) = 2sX(s) + 2X(s) + X(s)$$



TARUN KAUSHIK

(14.)

$$\Rightarrow (s^2 - 2s + 1) X(s) = \frac{1}{s-1} + 4s - 5$$

$$\Rightarrow X(s) = \frac{1}{(s-1)^3} + \frac{4s}{(s-1)^2} - \frac{5}{(s-1)^2}$$

$$= \frac{1}{(s-1)^3} + 4 \left( \frac{s+1-1}{(s-1)^2} \right) - \frac{5}{(s-1)^2}$$

$$= \frac{1}{(s-1)^3} + 4 \left( \frac{s-1}{(s-1)^2} + \frac{1}{(s-1)^2} \right) - \frac{5}{(s-1)^2}$$

$$= \frac{1}{(s-1)^3} + \frac{4}{(s-1)} + \frac{4}{(s-1)^2} - \frac{5}{(s-1)^2}$$

$$\Rightarrow X(s) = \frac{1}{(s-1)^3} + \frac{4}{(s-1)} - \frac{1}{(s-1)^2}$$

Now Take I.L.T.

$$x(t) = e^t \frac{t^2}{2} + 4e^t - te^t$$

$$\boxed{x(t) = e^t \left( \frac{t^2}{2} - t + 4 \right)} \text{ Ans.}$$



## # Laplace Transform of Periodic Function :-

Let  $f(t)$  be a periodic function with period  $T$  then  $L[f(t)]$  is defined as

$$L[f(t)] = \frac{1}{1-e^{-sT}} \int_0^T e^{-st} f(t) dt$$

Que:-  $f(t) = \begin{cases} 3, & 0 < t < 2 \\ 0, & 2 < t < 4 \end{cases}$

$$f(t+4) = f(t)$$

Find  $L[f(t)]$ .

Sol.

$$L[f(t)] = \frac{1}{1-e^{-4s}} \left[ \int_0^2 e^{-st} (3) dt + \int_2^4 e^{-st} (0) dt \right]$$

Formula :-


$$(1) L[t^n] = \frac{\sqrt{n+1}}{s^{n+1}} = \begin{cases} \frac{n!}{s^{n+1}} & ; n \in \mathbb{N} \\ \frac{n\sqrt{n}}{s^{n+1}} & ; n > 0 \end{cases}$$



TARUN KAUSHIK



**+91-9416421592**



**TARUN KAUSHIK**