

ASSIGNMENT 3

HARITHA R
AI20BTECH11010

Download all python codes from

<https://github.com/harithar1234/EE3900-Haritha/blob/main/assignment3/assignment3.py>

QUESTION

Construction/Q 2.18

Draw a circle with centre B and radius 6. If C be a point 10 units away from its centre, construct the pair of tangents AC and CD to the circle.

SOLUTION

Theorem 0.1. *The points of intersection of the line*

$$L: \mathbf{x} = \mathbf{q} + \mu \mathbf{m} \quad \mu \in \mathbb{R} \quad (0.0.1)$$

with a general conic are given by

$$\mathbf{x}_i = \mathbf{q} + \mu_i \mathbf{m} \quad (0.0.2)$$

where

$$\mu_i = \frac{1}{\mathbf{m}^T \mathbf{V} \mathbf{m}} \left(-\mathbf{m}^T (\mathbf{V} \mathbf{q} + \mathbf{u}) \pm \sqrt{[\mathbf{m}^T (\mathbf{V} \mathbf{q} + \mathbf{u})]^2 - (\mathbf{q}^T \mathbf{V} \mathbf{q} + 2\mathbf{u}^T \mathbf{q} + f)(\mathbf{m}^T \mathbf{V} \mathbf{m})} \right) \quad (0.0.3)$$

The general form of a conic is given by

$$\mathbf{x}^T \mathbf{V} \mathbf{x} + 2\mathbf{u}^T \mathbf{x} + f = 0 \quad (0.0.4)$$

for a circle, $\mathbf{V} = \mathbf{I}$ and $\mathbf{u} = -\mathbf{B}$ where \mathbf{B} is the center of the circle.

Let \mathbf{q} be the locus of the point of tangency from point \mathbf{C} , the distance of \mathbf{C} from \mathbf{B} is d

$$(\mathbf{q} - \mathbf{B})^T (\mathbf{q} - \mathbf{C}) = 0 \quad (0.0.5)$$

$$(\mathbf{q} + \mathbf{u})^T (\mathbf{q} - \mathbf{C}) = 0 \quad (0.0.6)$$

$$\mathbf{q}^T \mathbf{q} + \mathbf{u}^T \mathbf{q} - \mathbf{u}^T \mathbf{C} - \mathbf{q}^T \mathbf{C} = 0 \quad (0.0.7)$$

Using (0.0.4)

$$(\mathbf{u} + \mathbf{C})^T \mathbf{q} = -f - \mathbf{u}^T \mathbf{C} \quad (0.0.8)$$

Let $\mathbf{n} = \mathbf{u} + \mathbf{C}$ and $c = -f - \mathbf{u}^T \mathbf{C}$ This is equation of a line, let $\mathbf{q} = \mathbf{a}$ be a point that lies on this line

$$\therefore \mathbf{q} = \mathbf{a} + \lambda \begin{pmatrix} -\mathbf{e}_1^T \mathbf{n} \\ \mathbf{e}_2^T \mathbf{n} \end{pmatrix} \quad (0.0.9)$$

We need to find the intersection point of this with the given circle.

Using (0.0.3)

$$\mathbf{q} = \mathbf{a} + \mu_i \mathbf{m} \quad (0.0.10)$$

$$\mu_i = \frac{1}{d^2} \left(-\mathbf{m}^T (\mathbf{a} + \mathbf{u}) \pm \sqrt{[\mathbf{m}^T (\mathbf{a} + \mathbf{u})]^2 - (\mathbf{a}^T \mathbf{a} + 2\mathbf{u}^T \mathbf{a} + f)d^2} \right) \quad (0.0.11)$$

Now, we can confirm the solution by checking for $\mathbf{u} = \mathbf{0}$, $\mathbf{C} = d\mathbf{e}_1$, $f = -r^2$

$$\implies \mathbf{n} = d\mathbf{e}_1, \mathbf{m} = d\mathbf{e}_2 \quad (0.0.12)$$

An arbitrary choice of \mathbf{a} could be $\frac{r^2}{d}\mathbf{e}_1$,

$$\mu_i = \pm \frac{1}{d^2} \sqrt{(r^2 d^2 - r^4)} \quad (0.0.13)$$

$$\mathbf{q} = \frac{r^2}{d}\mathbf{e}_1 \pm r \sqrt{1 - \frac{r^2}{d^2}} \mathbf{e}_2 \quad (0.0.14)$$

\mathbf{A}, \mathbf{D} are the corresponding points of tangency from $\mathbf{C} = 10\mathbf{e}_1$

Using (0.0.14), we obtain all the points of tangency. $d=10$ and $r=6$.

Therefore, $\mathbf{A} = \begin{pmatrix} 3.6 \\ 4.8 \end{pmatrix}$ and $\mathbf{D} = \begin{pmatrix} 3.6 \\ -4.8 \end{pmatrix}$




Fig. 1: Plot of the tangents