

Polychaeta

Species	Rated		Total		In top 5		Feeding type	Motility	Mouth-part
	%		%		%				
<i>Scoloplos armiger</i>	151	16,0	1434	8,3	45	14,3	S ¹	M ¹	X ¹
<i>Chaetozone setosa</i>	99	10,5	1406	8,1	23	7,3	S ¹	D ¹ /M ¹	T ¹
<i>Mediomastus fragilis</i>	98	10,4	1753	10,2	25	7,9	S ¹	M ¹	X ¹
<i>Pholoe inornata</i>	43	4,4	480	2,8	14	4,4	C ¹	M ¹	J ¹
<i>Cirratulus cirratus</i>	37	2,9	394	2,3	9	2,9	S ¹	D ¹ /M ¹	T ¹
<i>Praxillella praetermissa</i>	31	3,3	447	2,6	11	3,5	S ¹	S ¹	X ¹
<i>Aricidea suecica</i>	25	2,6	793	4,6	6	1,9	H ¹ /S ¹	M ¹	X ¹
<i>Capitella capitata</i>	22	2,3	285	1,7	11	3,5	S ¹	M ¹	X ¹
<i>Spio filicornis</i>	20	2,1	820	4,8	7	2,2	F ¹ /S ¹	D ¹ /M ¹	T ¹
<i>Scalibregma inflatum</i>	20	2,1	381	2,2	9	2,1	S ¹	M ¹	X ¹
<i>Exogone naidina</i>	19	1,9	876	5,1	6	1,9	H ¹ /C ¹	M ¹	J ¹
<i>Levinsonia gracilis</i>	16	1,7	294	1,7	9	2,9	H ¹ /S ¹	M ¹	X ¹
<i>Nephtys caeca</i>	15	1,6	35	0,2	5	1,6	C ¹	M ¹	J ¹
<i>Pectinaria koreni</i>	14	1,5	167	1,0	7	2,2	S ¹	M ¹	X ¹
<i>Nephtys hombergi</i>	14	1,5	218	1,3	5	1,6	C ¹	M ¹	J ¹
<i>Polydora ciliata</i>	14	1,5	1131	6,6	4	1,3	F ¹ /S ¹	D ¹ /M ¹	T ¹
<i>Dipolydora quadrilobata</i>	12	1,3	460	2,7	5	1,6	F ¹ /S ¹	D ¹ /M ¹	T ¹
<i>Ophelina acuminata</i>	11	1,2	179	1,0	5	1,6	S ¹	M ¹	X ¹
<i>Lanice conchilega</i>	10	1,1	83	0,5	2	0,6	S ¹	S ¹	T ¹
<i>Goniada maculata</i>	9	1,0	140	0,8	4	1,3	C ¹	M ⁶	J ¹
<i>Nephtys ciliata</i>	8	0,8	133	0,8	5	1,6	C ¹	M ¹	J ¹
<i>Pseudopolydora pulchra</i>	8	0,8	155	0,9	4	1,3	F ¹ /S ¹	D ¹ /M ¹	T ¹
<i>Syllidae</i> sp, T187	7	0,7	141	0,8	5	1,6	C ¹	M ¹	J ¹
<i>Eumida sanguinea</i>	7	0,7	60	0,3	2	0,6	S ¹ /C ¹	M ¹	X ¹
<i>Myriochele oculata</i>	6	0,6	159	0,9	3	1,0	S ¹	D ¹	T ¹
<i>Ophryotrocha hartmanni</i>	6	0,6	150	0,9	3	1,0	H ¹ /C ¹	M ¹	J ¹
<i>Anobothrus gracilis</i>	5	0,5	41	0,2	1	0,3	S ¹	S ¹	T ¹
<i>Diplocirrus glaucus</i>	4	0,4	142	0,8	2	0,6	F ¹ /S ¹	D ¹ /M ¹	T ¹
<i>Chone fauveli</i>	4	0,4	10	0,1	1	0,3	F ¹	S ¹	T ¹
<i>Exogone verugera</i>	3	0,3	45	0,3	2	0,6	H ¹ /C ¹	M ¹	J ¹
<i>Pherusa flabellata</i>	3	0,3	32	0,2	1	0,3	F ¹ /S ¹	D ¹ /M ¹	T ¹
<i>Maldanidae</i> sp a	3	0,3	23	0,1	1	0,3	S ¹	S ¹	X ¹
<i>Eteone longa</i>	3	0,3	108	0,6	1	0,3	S ² /C ²	M ²	X ²
<i>Owenidae</i> sp	2	0,2	9	0,1	1	0,3	F ¹ /S ¹	D ¹	T ¹
<i>Cirratulidae</i> sp	2	0,2	14	0,1	1	0,3	S ¹	D ¹ /M ¹	T ¹
<i>Dipolydora socialis</i>	1	0,1	38	0,2	1	0,3	F ¹ /S ¹	D ¹ /M ¹	T ¹
<i>Travisia forbesii</i>	1	0,1	4	0,0	1	0,3	S ¹	M ¹	X ¹
<i>Kefersteinia cirrata</i>	1	0,1	38	0,2	1	0,3	C ¹	M ¹	J ¹
<i>Rhodine gracilior</i>	1	0,1	45	0,3	1	0,3	S ¹	S ¹	X ¹

Table 4a. Top five polychaetes from all samples showed as a rated value, total abundance and how often the species has been among the top five species. Information on feeding type, motility and mouthpart is also shown.

Feeding type: H: Herbivore. C: Carnivore. F: Filter-feeding/Suspension feeding S: Surface deposit feeding.
Motility: M: Motile. D: Discretely motile. S: Sessile.

Mouthparts: J: Jawed. X: Non-jawed. T: Tentaculate. P: Pumping.