NumPy Practice Assignment

- 1. Create an array of 10 fives
- 2. Create an array of the integers from 10 to 50
- 3. Create an array of all the even integers from 10 to 50
- 4. Create a 3x3 matrix with values ranging from 0 to 8
- 5. Use NumPy to generate an array of 25 random numbers sampled from a standard normal distribution
- 6. Create the following array:

```
array([0. , 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.11, 0.12, 0.13, 0.14, 0.15, 0.16, 0.17, 0.18, 0.19, 0.2, 0.21, 0.22, 0.23, 0.24, 0.25, 0.26, 0.27, 0.28, 0.29, 0.3, 0.31, 0.32, 0.33, 0.34, 0.35, 0.36, 0.37, 0.38, 0.39, 0.4, 0.41, 0.42, 0.43, 0.44, 0.45, 0.46, 0.47, 0.48, 0.49, 0.5, 0.51, 0.52, 0.53, 0.54, 0.55, 0.56, 0.57, 0.58, 0.59, 0.6, 0.61, 0.62, 0.63, 0.64, 0.65, 0.66, 0.67, 0.68, 0.69, 0.7, 0.71, 0.72, 0.73, 0.74, 0.75, 0.76, 0.77, 0.78, 0.79, 0.8, 0.81, 0.82, 0.83, 0.84, 0.85, 0.86, 0.87, 0.88, 0.89, 0.9, 0.91, 0.92, 0.93, 0.94, 0.95, 0.96, 0.97, 0.98, 0.99, 1. ])
```

- 7. Create an array of 20 linearly spaced points between 0 and 1
- 8. Generate the following matrix, mat,

```
array([[ 1, 2, 3, 4, 5], [ 6, 7, 8, 9, 10], [11, 12, 13, 14, 15], [16, 17, 18, 19, 20], [21, 22, 23, 24, 25]])
```

and extract the following using the indexing and slicing:

```
array([[12, 13, 14, 15],

[17, 18, 19, 20],

[22, 23, 24, 25]])

array([[2],

[7],

[12]])
```

Get the sum of all the values in mat

Get the sum of all the rows and columns in mat