
Comparison of SAD variance among methods to adjust for population structure



7	5	5	5	7	8	Hair color	3.07	9.21	2.41	3.43	0.98	0.98
4	6	6	4	7	7	Skin color	8.88	3.49	3.45	4.24	4.77	2.17
4	4	2	4	5	6	Diastolic blood pressure	8.13	2.28	3.88	3.70	1.28	1.27
2	1	2	2	4	3	Height	5.08	1.96	5.03	1.99	1.33	6.67
3	3	3	3	2	1	Forced vital capacity	2.88	2.99	5.42	5.99	2.46	2.51
2	3	3	3	0	1	Years of schooling	2.48	5.42	2.11	1.84	1.28	1.22
2	1	1	1	2	3	Birth weight	5.06	3.17	3.08	2.78	2.91	2.43
1	0	1	0	2	3	Neuroticism score	1.57	4.65	3.26	1.54	4.40	1.35
0	1	1	0	2	2	Household income	3.53	1.34	1.35	1.32	0.94	0.95
1	1	1	1	0	2	Waist circumference	1.88	3.12	2.39	2.85	0.90	0.90
1	2	2	0	2	1	Overall health rating	2.05	2.15	3.03	2.11	1.52	1.67
1	1	1	0	0	1	Alcohol intake frequency	2.36	2.35	2.92	2.23	2.40	2.34
0	1	1	0	0	0	Hand grip strength	1.82	2.86	1.72	1.83	1.35	1.84
1	1	1	1	1	0	Hip circumference	2.65	1.81	2.00	1.65	1.22	1.45
1	0	1	0	0	0	Pack years of smoking	2.29	2.14	1.94	1.91	2.09	2.17
1	1	1	1	0	1	Body mass index	2.20	1.33	1.34	1.30	1.20	1.14
0	0	0	0	1	1	Pulse rate	1.96	1.95	1.96	1.94	1.29	1.29