

Linearity: It adds up ...  
... but the converse doesn't hold.

Matt McDevitt

Scott Harper

SUMS Lunchtime Talk

21st April 2015

Linearity: It adds up ...  
... but the converse doesn't hold.

Matt McDevitt

Scott Harper

SUMS Lunchtime Talk

21st April 2015

# Non-Constructive Proofs

# Non-Constructive Proofs

## Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

# Non-Constructive Proofs

## Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

## Proof

Consider  $\sqrt{2}^{\sqrt{2}}$ .

# Non-Constructive Proofs

## Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

## Proof

Consider  $\sqrt{2}^{\sqrt{2}}$ .

**Case 1:**  $\sqrt{2}^{\sqrt{2}} \in \mathbb{Q}$

# Non-Constructive Proofs

## Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

## Proof

Consider  $\sqrt{2}^{\sqrt{2}}$ .

**Case 1:**  $\sqrt{2}^{\sqrt{2}} \in \mathbb{Q}$     We are done!  $\sqrt{2}^{\sqrt{2}}$  is rational.

# Non-Constructive Proofs

## Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

## Proof

Consider  $\sqrt{2}^{\sqrt{2}}$ .

**Case 1:**  $\sqrt{2}^{\sqrt{2}} \in \mathbb{Q}$     We are done!  $\sqrt{2}^{\sqrt{2}}$  is rational.

**Case 2:**  $\sqrt{2}^{\sqrt{2}} \notin \mathbb{Q}$

# Non-Constructive Proofs

## Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

## Proof

Consider  $\sqrt{2}^{\sqrt{2}}$ .

**Case 1:**  $\sqrt{2}^{\sqrt{2}} \in \mathbb{Q}$     We are done!  $\sqrt{2}^{\sqrt{2}}$  is rational.

**Case 2:**  $\sqrt{2}^{\sqrt{2}} \notin \mathbb{Q}$      $\left(\sqrt{2}^{\sqrt{2}}\right)^{\sqrt{2}}$

# Non-Constructive Proofs

## Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

## Proof

Consider  $\sqrt{2}^{\sqrt{2}}$ .

**Case 1:**  $\sqrt{2}^{\sqrt{2}} \in \mathbb{Q}$     We are done!  $\sqrt{2}^{\sqrt{2}}$  is rational.

**Case 2:**  $\sqrt{2}^{\sqrt{2}} \notin \mathbb{Q}$      $\left(\sqrt{2}^{\sqrt{2}}\right)^{\sqrt{2}} = \sqrt{2}^{\sqrt{2}\sqrt{2}}$

# Non-Constructive Proofs

## Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

## Proof

Consider  $\sqrt{2}^{\sqrt{2}}$ .

**Case 1:**  $\sqrt{2}^{\sqrt{2}} \in \mathbb{Q}$     We are done!  $\sqrt{2}^{\sqrt{2}}$  is rational.

**Case 2:**  $\sqrt{2}^{\sqrt{2}} \notin \mathbb{Q}$      $\left(\sqrt{2}^{\sqrt{2}}\right)^{\sqrt{2}} = \sqrt{2}^{\sqrt{2}\sqrt{2}} = \sqrt{2}^2$

# Non-Constructive Proofs

## Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

## Proof

Consider  $\sqrt{2}^{\sqrt{2}}$ .

**Case 1:**  $\sqrt{2}^{\sqrt{2}} \in \mathbb{Q}$     We are done!  $\sqrt{2}^{\sqrt{2}}$  is rational.

**Case 2:**  $\sqrt{2}^{\sqrt{2}} \notin \mathbb{Q}$      $\left(\sqrt{2}^{\sqrt{2}}\right)^{\sqrt{2}} = \sqrt{2}^{\sqrt{2}\sqrt{2}} = \sqrt{2}^2 = 2 \in \mathbb{Q}$ .

# Non-Constructive Proofs

## Theorem

*There exist irrational numbers  $a$  and  $b$  such that  $a^b$  is rational.*

## Proof

Consider  $\sqrt{2}^{\sqrt{2}}$ .

**Case 1:**  $\sqrt{2}^{\sqrt{2}} \in \mathbb{Q}$     We are done!  $\sqrt{2}^{\sqrt{2}}$  is rational.

**Case 2:**  $\sqrt{2}^{\sqrt{2}} \notin \mathbb{Q}$      $\left(\sqrt{2}^{\sqrt{2}}\right)^{\sqrt{2}} = \sqrt{2}^{\sqrt{2}\sqrt{2}} = \sqrt{2}^2 = 2 \in \mathbb{Q}$ . ■

# Real Numbers and Rational Numbers

# Real Numbers and Rational Numbers

Let  $x, y, z \in \mathbb{R}$  and let  $q, r \in \mathbb{Q}$ .

# Real Numbers and Rational Numbers

Let  $x, y, z \in \mathbb{R}$  and let  $q, r \in \mathbb{Q}$ .

(i)  $(x + y) + z = x + (y + z)$

# Real Numbers and Rational Numbers

Let  $x, y, z \in \mathbb{R}$  and let  $q, r \in \mathbb{Q}$ .

- (i)  $(x + y) + z = x + (y + z)$
- (ii)  $x + 0 = 0 + x = x$

# Real Numbers and Rational Numbers

Let  $x, y, z \in \mathbb{R}$  and let  $q, r \in \mathbb{Q}$ .

- (i)  $(x + y) + z = x + (y + z)$
- (ii)  $x + 0 = 0 + x = x$
- (iii)  $x + (-x) = 0 = (-x) + x$

# Real Numbers and Rational Numbers

Let  $x, y, z \in \mathbb{R}$  and let  $q, r \in \mathbb{Q}$ .

- (i)  $(x + y) + z = x + (y + z)$
- (ii)  $x + 0 = 0 + x = x$
- (iii)  $x + (-x) = 0 = (-x) + x$
- (iv)  $x + y = y + x$

# Real Numbers and Rational Numbers

Let  $x, y, z \in \mathbb{R}$  and let  $q, r \in \mathbb{Q}$ .

- (i)  $(x + y) + z = x + (y + z)$
- (ii)  $x + 0 = 0 + x = x$
- (iii)  $x + (-x) = 0 = (-x) + x$
- (iv)  $x + y = y + x$
- (v)  $(qr)x = q(rx)$

# Real Numbers and Rational Numbers

Let  $x, y, z \in \mathbb{R}$  and let  $q, r \in \mathbb{Q}$ .

- (i)  $(x + y) + z = x + (y + z)$
- (ii)  $x + 0 = 0 + x = x$
- (iii)  $x + (-x) = 0 = (-x) + x$
- (iv)  $x + y = y + x$
- (v)  $(qr)x = q(rx)$
- (vi)  $1x = x$

# Real Numbers and Rational Numbers

Let  $x, y, z \in \mathbb{R}$  and let  $q, r \in \mathbb{Q}$ .

- (i)  $(x + y) + z = x + (y + z)$
- (ii)  $x + 0 = 0 + x = x$
- (iii)  $x + (-x) = 0 = (-x) + x$
- (iv)  $x + y = y + x$
- (v)  $(qr)x = q(rx)$
- (vi)  $1x = x$
- (vii)  $q(x + y) = qx + qr$

# Real Numbers and Rational Numbers

Let  $x, y, z \in \mathbb{R}$  and let  $q, r \in \mathbb{Q}$ .

- (i)  $(x + y) + z = x + (y + z)$
- (ii)  $x + 0 = 0 + x = x$
- (iii)  $x + (-x) = 0 = (-x) + x$
- (iv)  $x + y = y + x$
- (v)  $(qr)x = q(rx)$
- (vi)  $1x = x$
- (vii)  $q(x + y) = qx + qr$
- (viii)  $(q + r)x = qx + rx.$

# Real Numbers and Rational Numbers

Let  $x, y, z \in \mathbb{R}$  and let  $q, r \in \mathbb{Q}$ .

- (i)  $(x + y) + z = x + (y + z)$
- (ii)  $x + 0 = 0 + x = x$
- (iii)  $x + (-x) = 0 = (-x) + x$
- (iv)  $x + y = y + x$
- (v)  $(qr)x = q(rx)$
- (vi)  $1x = x$
- (vii)  $q(x + y) = qx + qr$
- (viii)  $(q + r)x = qx + rx.$

## Theorem

$\mathbb{R}$  is a vector space over  $\mathbb{Q}$ .

# Bases

Let  $V$  be a vector space over a field  $F$ .

# Bases

Let  $V$  be a vector space over a field  $F$ .

## Definition (Familiar)

A set  $\mathcal{B} = \{x_1, \dots, x_n\} \subseteq V$  is a **basis** for  $V$  if:

# Bases

Let  $V$  be a vector space over a field  $F$ .

## Definition (Familiar)

A set  $\mathcal{B} = \{x_1, \dots, x_n\} \subseteq V$  is a **basis** for  $V$  if:

**Spanning** For every  $x \in V$  there exist  $q_1, \dots, q_n \in F$  such that  
$$x = q_1x_1 + \cdots + q_nx_n.$$

# Bases

Let  $V$  be a vector space over a field  $F$ .

## Definition (Familiar)

A set  $\mathcal{B} = \{x_1, \dots, x_n\} \subseteq V$  is a **basis** for  $V$  if:

**Spanning** For every  $x \in V$  there exist  $q_1, \dots, q_n \in F$  such that  
$$x = q_1x_1 + \dots + q_nx_n.$$

**Linear Independence** If  $q_1x_1 + \dots + q_nx_n = 0$  then  $q_1 = \dots = q_n = 0$ .

# Bases

Let  $V$  be a vector space over a field  $F$ .

## Definition (Familiar)

A set  $\mathcal{B} = \{x_1, \dots, x_n\} \subseteq V$  is a **basis** for  $V$  if:

**Spanning** For every  $x \in V$  there exist  $q_1, \dots, q_n \in F$  such that

$$x = q_1x_1 + \cdots + q_nx_n.$$

**Linear Independence** If  $q_1x_1 + \cdots + q_nx_n = 0$  then  $q_1 = \cdots = q_n = 0$ .

## Definition (New)

A set  $\mathcal{B} \subseteq V$  is a **basis** for  $V$  if:

# Bases

Let  $V$  be a vector space over a field  $F$ .

## Definition (Familiar)

A set  $\mathcal{B} = \{x_1, \dots, x_n\} \subseteq V$  is a **basis** for  $V$  if:

**Spanning** For every  $x \in V$  there exist  $q_1, \dots, q_n \in F$  such that  
$$x = q_1x_1 + \dots + q_nx_n.$$

**Linear Independence** If  $q_1x_1 + \dots + q_nx_n = 0$  then  $q_1 = \dots = q_n = 0$ .

## Definition (New)

A set  $\mathcal{B} \subseteq V$  is a **basis** for  $V$  if:

**Spanning** For each  $x \in V$  there exist  $x_1, \dots, x_n \in \mathcal{B}$  and  
 $q_1, \dots, q_n \in F$  such that  $x = q_1x_1 + \dots + q_nx_n,$

# Bases

Let  $V$  be a vector space over a field  $F$ .

## Definition (Familiar)

A set  $\mathcal{B} = \{x_1, \dots, x_n\} \subseteq V$  is a **basis** for  $V$  if:

**Spanning** For every  $x \in V$  there exist  $q_1, \dots, q_n \in F$  such that  
$$x = q_1x_1 + \dots + q_nx_n.$$

**Linear Independence** If  $q_1x_1 + \dots + q_nx_n = 0$  then  $q_1 = \dots = q_n = 0$ .

## Definition (New)

A set  $\mathcal{B} \subseteq V$  is a **basis** for  $V$  if:

**Spanning** For each  $x \in V$  there exist  $x_1, \dots, x_n \in \mathcal{B}$  and  
 $q_1, \dots, q_n \in F$  such that  $x = q_1x_1 + \dots + q_nx_n,$

**Linear Independence** For any  $x_1, \dots, x_m \in \mathcal{B}$ , if  $q_1x_1 + \dots + q_mx_m = 0$   
then  $q_1 = \dots = q_m = 0$ .

# Why do we care?

# Why do we care?

Let  $\mathcal{B}$  be a basis for  $\mathbb{R}$  over  $\mathbb{Q}$ .

# Why do we care?

Let  $\mathcal{B}$  be a basis for  $\mathbb{R}$  over  $\mathbb{Q}$ . Fix  $b \in \mathcal{B}$ .

# Why do we care?

Let  $\mathcal{B}$  be a basis for  $\mathbb{R}$  over  $\mathbb{Q}$ . Fix  $b \in \mathcal{B}$ .

Define  $f : \mathcal{B} \rightarrow \mathbb{R}$  as

$$f(x) = \begin{cases} x + 1 & \text{if } x = b \\ x & \text{otherwise} \end{cases}$$

# Why do we care?

Let  $\mathcal{B}$  be a basis for  $\mathbb{R}$  over  $\mathbb{Q}$ . Fix  $b \in \mathcal{B}$ .

Define  $f : \mathcal{B} \rightarrow \mathbb{R}$  as

$$f(x) = \begin{cases} x + 1 & \text{if } x = b \\ x & \text{otherwise} \end{cases}$$

For each  $x \in \mathbb{R}$  write  $x = q_1 b_1 + \cdots + q_n b_n$

# Why do we care?

Let  $\mathcal{B}$  be a basis for  $\mathbb{R}$  over  $\mathbb{Q}$ . Fix  $b \in \mathcal{B}$ .

Define  $f : \mathcal{B} \rightarrow \mathbb{R}$  as

$$f(x) = \begin{cases} x + 1 & \text{if } x = b \\ x & \text{otherwise} \end{cases}$$

For each  $x \in \mathbb{R}$  write  $x = q_1 b_1 + \cdots + q_n b_n$  and define

$$f^*(x) = q_1 f(b_1) + \cdots + q_n f(b_n).$$

# Why do we care?

Let  $\mathcal{B}$  be a basis for  $\mathbb{R}$  over  $\mathbb{Q}$ . Fix  $b \in \mathcal{B}$ .

Define  $f : \mathcal{B} \rightarrow \mathbb{R}$  as

$$f(x) = \begin{cases} x + 1 & \text{if } x = b \\ x & \text{otherwise} \end{cases}$$

For each  $x \in \mathbb{R}$  write  $x = q_1 b_1 + \cdots + q_n b_n$  and define

$$f^*(x) = q_1 f(b_1) + \cdots + q_n f(b_n).$$

**Claim:**  $f^*$  is non-linear.

# Why do we care?

Let  $\mathcal{B}$  be a basis for  $\mathbb{R}$  over  $\mathbb{Q}$ . Fix  $b \in \mathcal{B}$ .

Define  $f : \mathcal{B} \rightarrow \mathbb{R}$  as

$$f(x) = \begin{cases} x + 1 & \text{if } x = b \\ x & \text{otherwise} \end{cases}$$

For each  $x \in \mathbb{R}$  write  $x = q_1 b_1 + \cdots + q_n b_n$  and define

$$f^*(x) = q_1 f(b_1) + \cdots + q_n f(b_n).$$

**Claim:**  $f^*$  is non-linear.  $f^*$  is additive.

# Why do we care?

Let  $\mathcal{B}$  be a basis for  $\mathbb{R}$  over  $\mathbb{Q}$ . Fix  $b \in \mathcal{B}$ .

Define  $f : \mathcal{B} \rightarrow \mathbb{R}$  as

$$f(x) = \begin{cases} x + 1 & \text{if } x = b \\ x & \text{otherwise} \end{cases}$$

For each  $x \in \mathbb{R}$  write  $x = q_1 b_1 + \cdots + q_n b_n$  and define

$$f^*(x) = q_1 f(b_1) + \cdots + q_n f(b_n).$$

**Claim:**  $f^*$  is non-linear.  $f^*$  is additive.

## Theorem

*Every vector space has a basis.*

## Why do we care?

Let  $\mathcal{B}$  be a basis for  $\mathbb{R}$  over  $\mathbb{Q}$ . Fix  $b \in \mathcal{B}$ .

Define  $f : \mathcal{B} \rightarrow \mathbb{R}$  as

$$f(x) = \begin{cases} x + 1 & \text{if } x = b \\ x & \text{otherwise} \end{cases}$$

For each  $x \in \mathbb{R}$  write  $x = q_1 b_1 + \cdots + q_n b_n$  and define

$$f^*(x) = q_1 f(b_1) + \cdots + q_n f(b_n).$$

**Claim:**  $f^*$  is non-linear.  $f^*$  is additive.

### Theorem

*Every vector space has a basis.*

### Corollary

*There exists an additive function  $f : \mathbb{R} \rightarrow \mathbb{R}$  which is non-linear.*

# Every vector space has a basis

# Every vector space has a basis

## Example 1 ( $\mathbb{R}^3$ )

# Every vector space has a basis

Example 1 ( $\mathbb{R}^3$ )

$$\mathcal{B}_1 = \{(1, 0, 0)\}$$

# Every vector space has a basis

## Example 1 ( $\mathbb{R}^3$ )

$$\mathcal{B}_1 = \{(1, 0, 0)\} \quad \mathcal{B}_2 = \{(1, 0, 0), (0, 1, 0)\}$$

# Every vector space has a basis

## Example 1 ( $\mathbb{R}^3$ )

$$\mathcal{B}_1 = \{(1, 0, 0)\} \quad \mathcal{B}_2 = \{(1, 0, 0), (0, 1, 0)\} \quad \mathcal{B}_3 = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$$

# Every vector space has a basis

## Example 1 ( $\mathbb{R}^3$ )

$$\mathcal{B}_1 = \{(1, 0, 0)\} \quad \mathcal{B}_2 = \{(1, 0, 0), (0, 1, 0)\} \quad \mathcal{B}_3 = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$$

## Claim 1

*A maximal linearly independent set is a basis.*

# Every vector space has a basis

## Example 1 ( $\mathbb{R}^3$ )

$$\mathcal{B}_1 = \{(1, 0, 0)\} \quad \mathcal{B}_2 = \{(1, 0, 0), (0, 1, 0)\} \quad \mathcal{B}_3 = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$$

## Claim 1

*A maximal linearly independent set is a basis.*

## Proof

# Every vector space has a basis

## Example 1 ( $\mathbb{R}^3$ )

$$\mathcal{B}_1 = \{(1, 0, 0)\} \quad \mathcal{B}_2 = \{(1, 0, 0), (0, 1, 0)\} \quad \mathcal{B}_3 = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$$

## Claim 1

*A maximal linearly independent set is a basis.*

## Proof

Let  $\mathcal{B}$  be a maximal linearly independent set. Does  $\mathcal{B}$  span  $V$ ?

# Every vector space has a basis

## Example 1 ( $\mathbb{R}^3$ )

$$\mathcal{B}_1 = \{(1, 0, 0)\} \quad \mathcal{B}_2 = \{(1, 0, 0), (0, 1, 0)\} \quad \mathcal{B}_3 = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$$

## Claim 1

*A maximal linearly independent set is a basis.*

## Proof

Let  $\mathcal{B}$  be a maximal linearly independent set. Does  $\mathcal{B}$  span  $V$ ?

Certainly  $\mathcal{B}$  spans  $\mathcal{B}$ , so let  $x \notin \mathcal{B}$ .

# Every vector space has a basis

## Example 1 ( $\mathbb{R}^3$ )

$$\mathcal{B}_1 = \{(1, 0, 0)\} \quad \mathcal{B}_2 = \{(1, 0, 0), (0, 1, 0)\} \quad \mathcal{B}_3 = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$$

## Claim 1

*A maximal linearly independent set is a basis.*

## Proof

Let  $\mathcal{B}$  be a maximal linearly independent set. Does  $\mathcal{B}$  span  $V$ ?

Certainly  $\mathcal{B}$  spans  $\mathcal{B}$ , so let  $x \notin \mathcal{B}$ .

Since  $\mathcal{B}$  is a maximal l.i. set  $\mathcal{B} \cup \{x\}$  is not l.i.

# Every vector space has a basis

## Example 1 ( $\mathbb{R}^3$ )

$$\mathcal{B}_1 = \{(1, 0, 0)\} \quad \mathcal{B}_2 = \{(1, 0, 0), (0, 1, 0)\} \quad \mathcal{B}_3 = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$$

## Claim 1

*A maximal linearly independent set is a basis.*

## Proof

Let  $\mathcal{B}$  be a maximal linearly independent set. Does  $\mathcal{B}$  span  $V$ ?

Certainly  $\mathcal{B}$  spans  $\mathcal{B}$ , so let  $x \notin \mathcal{B}$ .

Since  $\mathcal{B}$  is a maximal l.i. set  $\mathcal{B} \cup \{x\}$  is not l.i.

There are  $x_1, \dots, x_n \in \mathcal{B} \cup \{x\}$  and  $q_1, \dots, q_n \in F$  s.t.

$$q_1x_1 + \dots + q_nx_n = 0.$$

# Every vector space has a basis

## Example 1 ( $\mathbb{R}^3$ )

$$\mathcal{B}_1 = \{(1, 0, 0)\} \quad \mathcal{B}_2 = \{(1, 0, 0), (0, 1, 0)\} \quad \mathcal{B}_3 = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$$

## Claim 1

*A maximal linearly independent set is a basis.*

## Proof

Let  $\mathcal{B}$  be a maximal linearly independent set. Does  $\mathcal{B}$  span  $V$ ?

Certainly  $\mathcal{B}$  spans  $\mathcal{B}$ , so let  $x \notin \mathcal{B}$ .

Since  $\mathcal{B}$  is a maximal l.i. set  $\mathcal{B} \cup \{x\}$  is not l.i.

There are  $x_1, \dots, x_n \in \mathcal{B} \cup \{x\}$  and  $q_1, \dots, q_n \in F$  s.t.

$$q_1x_1 + \dots + q_nx_n = 0.$$

Since  $\mathcal{B}$  is l.i. assume that  $x_1 = x$ .

# Every vector space has a basis

## Example 1 ( $\mathbb{R}^3$ )

$$\mathcal{B}_1 = \{(1, 0, 0)\} \quad \mathcal{B}_2 = \{(1, 0, 0), (0, 1, 0)\} \quad \mathcal{B}_3 = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$$

## Claim 1

*A maximal linearly independent set is a basis.*

## Proof

Let  $\mathcal{B}$  be a maximal linearly independent set. Does  $\mathcal{B}$  span  $V$ ?

Certainly  $\mathcal{B}$  spans  $\mathcal{B}$ , so let  $x \notin \mathcal{B}$ .

Since  $\mathcal{B}$  is a maximal l.i. set  $\mathcal{B} \cup \{x\}$  is not l.i.

There are  $x_1, \dots, x_n \in \mathcal{B} \cup \{x\}$  and  $q_1, \dots, q_n \in F$  s.t.

$$q_1x_1 + \dots + q_nx_n = 0.$$

Since  $\mathcal{B}$  is l.i. assume that  $x_1 = x$ .

$$\text{So } x = -\frac{q_2}{q_1}x_2 - \dots - \frac{q_n}{q_1}x_n.$$

# Every vector space has a basis

## Example 1 ( $\mathbb{R}^3$ )

$$\mathcal{B}_1 = \{(1, 0, 0)\} \quad \mathcal{B}_2 = \{(1, 0, 0), (0, 1, 0)\} \quad \mathcal{B}_3 = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$$

## Claim 1

*A maximal linearly independent set is a basis.*

## Proof

Let  $\mathcal{B}$  be a maximal linearly independent set. Does  $\mathcal{B}$  span  $V$ ?

Certainly  $\mathcal{B}$  spans  $\mathcal{B}$ , so let  $x \notin \mathcal{B}$ .

Since  $\mathcal{B}$  is a maximal l.i. set  $\mathcal{B} \cup \{x\}$  is not l.i.

There are  $x_1, \dots, x_n \in \mathcal{B} \cup \{x\}$  and  $q_1, \dots, q_n \in F$  s.t.

$$q_1x_1 + \dots + q_nx_n = 0.$$

Since  $\mathcal{B}$  is l.i. assume that  $x_1 = x$ .

So  $x = -\frac{q_2}{q_1}x_2 - \dots - \frac{q_n}{q_1}x_n$ . So  $\mathcal{B}$  spans  $V$ .



# Every vector space has a basis

# Every vector space has a basis

## Example 2 ( $\mathbb{R}[X]$ )

# Every vector space has a basis

## Example 2 ( $\mathbb{R}[X]$ )

$\mathcal{B}_1 = \{1\}$ ,

# Every vector space has a basis

## Example 2 ( $\mathbb{R}[X]$ )

$\mathcal{B}_1 = \{1\}$ ,  $\mathcal{B}_2 = \{1, X\}$ ,

# Every vector space has a basis

## Example 2 ( $\mathbb{R}[X]$ )

$\mathcal{B}_1 = \{1\}$ ,  $\mathcal{B}_2 = \{1, X\}$ ,  $\mathcal{B}_3 = \{1, X, X^2\}$ ,

# Every vector space has a basis

## Example 2 ( $\mathbb{R}[X]$ )

$\mathcal{B}_1 = \{1\}$ ,  $\mathcal{B}_2 = \{1, X\}$ ,  $\mathcal{B}_3 = \{1, X, X^2\}$ , ...

# Every vector space has a basis

## Example 2 ( $\mathbb{R}[X]$ )

$\mathcal{B}_1 = \{1\}, \mathcal{B}_2 = \{1, X\}, \mathcal{B}_3 = \{1, X, X^2\}, \dots$

$\mathcal{B} = \bigcup_{n \in \mathbb{N}} \mathcal{B}_n = \{1, X, X^2, X^3, \dots\}$

# Every vector space has a basis

## Example 2 ( $\mathbb{R}[X]$ )

$\mathcal{B}_1 = \{1\}$ ,  $\mathcal{B}_2 = \{1, X\}$ ,  $\mathcal{B}_3 = \{1, X, X^2\}$ , ...

$\mathcal{B} = \bigcup_{n \in \mathbb{N}} \mathcal{B}_n = \{1, X, X^2, X^3, \dots\}$

## Claim 2 – ε

Let  $X = \{\mathcal{B}_1, \mathcal{B}_2, \mathcal{B}_3, \dots\}$  be a collection of l.i. sets such that

$\mathcal{B}_1 \subseteq \mathcal{B}_2 \subseteq \mathcal{B}_3 \subseteq \dots$ . Then  $\mathcal{B} = \bigcup_{n \in \mathbb{N}} \mathcal{B}_n$  is l.i.

# Every vector space has a basis

## Example 2 ( $\mathbb{R}[X]$ )

$\mathcal{B}_1 = \{1\}$ ,  $\mathcal{B}_2 = \{1, X\}$ ,  $\mathcal{B}_3 = \{1, X, X^2\}$ , ...

$\mathcal{B} = \bigcup_{n \in \mathbb{N}} \mathcal{B}_n = \{1, X, X^2, X^3, \dots\}$

## Claim 2 – ε

Let  $X = \{\mathcal{B}_1, \mathcal{B}_2, \mathcal{B}_3, \dots\}$  be a collection of l.i. sets such that

$\mathcal{B}_1 \subseteq \mathcal{B}_2 \subseteq \mathcal{B}_3 \subseteq \dots$ . Then  $\mathcal{B} = \bigcup_{n \in \mathbb{N}} \mathcal{B}_n$  is l.i.

## Claim 2

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a collection of l.i. sets such that for any sets  $\mathcal{B}_i, \mathcal{B}_j \in X$ , either  $\mathcal{B}_i \subseteq \mathcal{B}_j$  or  $\mathcal{B}_j \subseteq \mathcal{B}_i$ . Then  $\mathcal{B} = \bigcup_{i \in I} \mathcal{B}_i$  is l.i.

# Every vector space has a basis

## Claim 2

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a collection of l.i. sets such that for any sets  $\mathcal{B}_i, \mathcal{B}_j \in X$ , either  $\mathcal{B}_i \subseteq \mathcal{B}_j$  or  $\mathcal{B}_j \subseteq \mathcal{B}_i$ . Then  $\mathcal{B} = \bigcup_{i \in I} \mathcal{B}_i$  is l.i.

# Every vector space has a basis

## Claim 2

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a collection of l.i. sets such that for any sets  $\mathcal{B}_i, \mathcal{B}_j \in X$ , either  $\mathcal{B}_i \subseteq \mathcal{B}_j$  or  $\mathcal{B}_j \subseteq \mathcal{B}_i$ . Then  $\mathcal{B} = \bigcup_{i \in I} \mathcal{B}_i$  is l.i.

## Proof

# Every vector space has a basis

## Claim 2

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a collection of l.i. sets such that for any sets  $\mathcal{B}_i, \mathcal{B}_j \in X$ , either  $\mathcal{B}_i \subseteq \mathcal{B}_j$  or  $\mathcal{B}_j \subseteq \mathcal{B}_i$ . Then  $\mathcal{B} = \bigcup_{i \in I} \mathcal{B}_i$  is l.i.

## Proof

Let  $x_1, \dots, x_n \in \mathcal{B}$  and  $q_1, \dots, q_n \in F$  be such that  $q_1x_1 + \dots + q_nx_n = 0$ .

# Every vector space has a basis

## Claim 2

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a collection of l.i. sets such that for any sets  $\mathcal{B}_i, \mathcal{B}_j \in X$ , either  $\mathcal{B}_i \subseteq \mathcal{B}_j$  or  $\mathcal{B}_j \subseteq \mathcal{B}_i$ . Then  $\mathcal{B} = \bigcup_{i \in I} \mathcal{B}_i$  is l.i.

## Proof

Let  $x_1, \dots, x_n \in \mathcal{B}$  and  $q_1, \dots, q_n \in F$  be such that  $q_1x_1 + \dots + q_nx_n = 0$ .

For  $i$  we know  $x_i \in \mathcal{B}_{j_i}$ .

# Every vector space has a basis

## Claim 2

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a collection of l.i. sets such that for any sets  $\mathcal{B}_i, \mathcal{B}_j \in X$ , either  $\mathcal{B}_i \subseteq \mathcal{B}_j$  or  $\mathcal{B}_j \subseteq \mathcal{B}_i$ . Then  $\mathcal{B} = \bigcup_{i \in I} \mathcal{B}_i$  is l.i.

## Proof

Let  $x_1, \dots, x_n \in \mathcal{B}$  and  $q_1, \dots, q_n \in F$  be such that  $q_1x_1 + \dots + q_nx_n = 0$ .

For  $i$  we know  $x_i \in \mathcal{B}_{j_i}$ .

Either  $\mathcal{B}_{j_1} \subseteq \mathcal{B}_{j_2}$  or  $\mathcal{B}_{j_2} \subseteq \mathcal{B}_{j_1}$ .

# Every vector space has a basis

## Claim 2

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a collection of l.i. sets such that for any sets  $\mathcal{B}_i, \mathcal{B}_j \in X$ , either  $\mathcal{B}_i \subseteq \mathcal{B}_j$  or  $\mathcal{B}_j \subseteq \mathcal{B}_i$ . Then  $\mathcal{B} = \bigcup_{i \in I} \mathcal{B}_i$  is l.i.

## Proof

Let  $x_1, \dots, x_n \in \mathcal{B}$  and  $q_1, \dots, q_n \in F$  be such that  $q_1x_1 + \dots + q_nx_n = 0$ .

For  $i$  we know  $x_i \in \mathcal{B}_{j_i}$ .

Either  $\mathcal{B}_{j_1} \subseteq \mathcal{B}_{j_2}$  or  $\mathcal{B}_{j_2} \subseteq \mathcal{B}_{j_1}$ . Assume that  $\mathcal{B}_{j_1} \subseteq \mathcal{B}_{j_2}$ .

# Every vector space has a basis

## Claim 2

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a collection of l.i. sets such that for any sets  $\mathcal{B}_i, \mathcal{B}_j \in X$ , either  $\mathcal{B}_i \subseteq \mathcal{B}_j$  or  $\mathcal{B}_j \subseteq \mathcal{B}_i$ . Then  $\mathcal{B} = \bigcup_{i \in I} \mathcal{B}_i$  is l.i.

## Proof

Let  $x_1, \dots, x_n \in \mathcal{B}$  and  $q_1, \dots, q_n \in F$  be such that  $q_1x_1 + \dots + q_nx_n = 0$ .

For  $i$  we know  $x_i \in \mathcal{B}_{j_i}$ .

Either  $\mathcal{B}_{j_1} \subseteq \mathcal{B}_{j_2}$  or  $\mathcal{B}_{j_2} \subseteq \mathcal{B}_{j_1}$ . Assume that  $\mathcal{B}_{j_1} \subseteq \mathcal{B}_{j_2}$ .

Then  $x_1, x_2 \in \mathcal{B}_{j_2}$ .

# Every vector space has a basis

## Claim 2

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a collection of l.i. sets such that for any sets  $\mathcal{B}_i, \mathcal{B}_j \in X$ , either  $\mathcal{B}_i \subseteq \mathcal{B}_j$  or  $\mathcal{B}_j \subseteq \mathcal{B}_i$ . Then  $\mathcal{B} = \bigcup_{i \in I} \mathcal{B}_i$  is l.i.

## Proof

Let  $x_1, \dots, x_n \in \mathcal{B}$  and  $q_1, \dots, q_n \in F$  be such that  $q_1x_1 + \dots + q_nx_n = 0$ .

For  $i$  we know  $x_i \in \mathcal{B}_{j_i}$ .

Either  $\mathcal{B}_{j_1} \subseteq \mathcal{B}_{j_2}$  or  $\mathcal{B}_{j_2} \subseteq \mathcal{B}_{j_1}$ . Assume that  $\mathcal{B}_{j_1} \subseteq \mathcal{B}_{j_2}$ .

Then  $x_1, x_2 \in \mathcal{B}_{j_2}$ .

Continuing this gives  $x_1, \dots, x_n \in \mathcal{B}_{j_n}$ .

# Every vector space has a basis

## Claim 2

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a collection of l.i. sets such that for any sets  $\mathcal{B}_i, \mathcal{B}_j \in X$ , either  $\mathcal{B}_i \subseteq \mathcal{B}_j$  or  $\mathcal{B}_j \subseteq \mathcal{B}_i$ . Then  $\mathcal{B} = \bigcup_{i \in I} \mathcal{B}_i$  is l.i.

## Proof

Let  $x_1, \dots, x_n \in \mathcal{B}$  and  $q_1, \dots, q_n \in F$  be such that  $q_1x_1 + \dots + q_nx_n = 0$ .

For  $i$  we know  $x_i \in \mathcal{B}_{j_i}$ .

Either  $\mathcal{B}_{j_1} \subseteq \mathcal{B}_{j_2}$  or  $\mathcal{B}_{j_2} \subseteq \mathcal{B}_{j_1}$ . Assume that  $\mathcal{B}_{j_1} \subseteq \mathcal{B}_{j_2}$ .

Then  $x_1, x_2 \in \mathcal{B}_{j_2}$ .

Continuing this gives  $x_1, \dots, x_n \in \mathcal{B}_{j_n}$ .

Since  $\mathcal{B}_{j_n}$  is linearly independent,  $q_1 = \dots = q_n = 0$ . ■

# Diversion: Set Theory

# Diversion: Set Theory

## Banach-Tarski Paradox

The unit sphere can be partitioned into finitely many parts which can be reassembled (following some isometries) to form two copies of the unit sphere.

# Diversion: Set Theory

## Banach-Tarski Paradox

The unit sphere can be partitioned into finitely many parts which can be reassembled (following some isometries) to form two copies of the unit sphere.

For any set  $\{A_i \mid i \in I\}$  of non-empty sets,  
there is a set  $X = \{a_i \mid i \in I\}$  such that  
 $a_i \in A_i$  for all  $i \in I$ .

# Diversion: Set Theory

## Banach-Tarski Paradox

The unit sphere can be partitioned into finitely many parts which can be reassembled (following some isometries) to form two copies of the unit sphere.

## Axiom of Choice

For any set  $\{A_i \mid i \in I\}$  of non-empty sets, there is a set  $X = \{a_i \mid i \in I\}$  such that  $a_i \in A_i$  for all  $i \in I$ .

# Diversion: Set Theory

## Banach-Tarski Paradox

The unit sphere can be partitioned into finitely many parts which can be reassembled (following some isometries) to form two copies of the unit sphere.

## Axiom of Choice

For any set  $\{A_i \mid i \in I\}$  of non-empty sets, there is a set  $X = \{a_i \mid i \in I\}$  such that  $a_i \in A_i$  for all  $i \in I$ .



(Source: <https://xkcd.com/804/>)

# Diversion: Set Theory

## Banach-Tarski Paradox

The unit sphere can be partitioned into finitely many parts which can be reassembled (following some isometries) to form two copies of the unit sphere.

## Axiom of Choice

For any set  $\{A_i \mid i \in I\}$  of non-empty sets, there is a set  $X = \{a_i \mid i \in I\}$  such that  $a_i \in A_i$  for all  $i \in I$ .

## Zorn's Lemma

Let  $X$  be a partially ordered set such that every totally ordered subset has an upper bound in  $X$ . Then  $X$  has a maximal element.



(Source: <https://xkcd.com/804/>)

# Diversion: Set Theory

# Diversion: Set Theory

## Zorn's Lemma

Let  $X$  be a partially ordered set such that every totally ordered subset has an upper bound in  $X$ . Then  $X$  has a maximal element.

# Diversion: Set Theory

## Zorn's Lemma

Let  $X$  be a partially ordered set such that every totally ordered subset has an upper bound in  $X$ . Then  $X$  has a maximal element.

## Definition

A **partial order** on a set  $X$  is a relation  $\leq$  such that for all  $x, y, z \in X$ :

- (i)  $x \leq x$ ;
- (ii) If  $x \leq y$  and  $y \leq x$  then  $x = y$ ;
- (iii) If  $x \leq y$  and  $y \leq z$  then  $x \leq z$ .

# Diversion: Set Theory

## Zorn's Lemma

Let  $X$  be a partially ordered set such that every totally ordered subset has an upper bound in  $X$ . Then  $X$  has a maximal element.

## Definition

A **partial order** on a set  $X$  is a relation  $\leq$  such that for all  $x, y, z \in X$ :

- (i)  $x \leq x$ ;
- (ii) If  $x \leq y$  and  $y \leq x$  then  $x = y$ ;
- (iii) If  $x \leq y$  and  $y \leq z$  then  $x \leq z$ .

## Definition

A **total order** on a set  $X$  is a partial order  $\leq$  such that for all  $x, y \in X$ ,

- (iv)  $x \leq y$  or  $y \leq x$ .

# Every vector space has a basis

# Every vector space has a basis

Let  $\mathcal{L}(V)$  be the set of all linearly independent subsets of  $V$  partially ordered by set containment.

# Every vector space has a basis

Let  $\mathcal{L}(V)$  be the set of all linearly independent subsets of  $V$  partially ordered by set containment.

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a totally ordered subset of  $\mathcal{L}(V)$ .

# Every vector space has a basis

Let  $\mathcal{L}(V)$  be the set of all linearly independent subsets of  $V$  partially ordered by set containment.

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a totally ordered subset of  $\mathcal{L}(V)$ . That is for any sets  $\mathcal{B}_i, \mathcal{B}_j \in X$ , either  $\mathcal{B}_i \subseteq \mathcal{B}_j$  or  $\mathcal{B}_j \subseteq \mathcal{B}_i$ .

# Every vector space has a basis

Let  $\mathcal{L}(V)$  be the set of all linearly independent subsets of  $V$  partially ordered by set containment.

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a totally ordered subset of  $\mathcal{L}(V)$ . That is for any sets  $\mathcal{B}_i, \mathcal{B}_j \in X$ , either  $\mathcal{B}_i \subseteq \mathcal{B}_j$  or  $\mathcal{B}_j \subseteq \mathcal{B}_i$ .

Let  $\mathcal{B} = \bigcup_{i \in I} \mathcal{B}_i$ .

# Every vector space has a basis

Let  $\mathcal{L}(V)$  be the set of all linearly independent subsets of  $V$  partially ordered by set containment.

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a totally ordered subset of  $\mathcal{L}(V)$ . That is for any sets  $\mathcal{B}_i, \mathcal{B}_j \in X$ , either  $\mathcal{B}_i \subseteq \mathcal{B}_j$  or  $\mathcal{B}_j \subseteq \mathcal{B}_i$ .

Let  $\mathcal{B} = \bigcup_{i \in I} \mathcal{B}_i$ . For all  $i \in I$ ,  $\mathcal{B}_i \subseteq \mathcal{B}$ .

# Every vector space has a basis

Let  $\mathcal{L}(V)$  be the set of all linearly independent subsets of  $V$  partially ordered by set containment.

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a totally ordered subset of  $\mathcal{L}(V)$ . That is for any sets  $\mathcal{B}_i, \mathcal{B}_j \in X$ , either  $\mathcal{B}_i \subseteq \mathcal{B}_j$  or  $\mathcal{B}_j \subseteq \mathcal{B}_i$ .

Let  $\mathcal{B} = \bigcup_{i \in I} \mathcal{B}_i$ . For all  $i \in I$ ,  $\mathcal{B}_i \subseteq \mathcal{B}$ . So  $\mathcal{B}$  is an upper bound for  $X$ .

# Every vector space has a basis

Let  $\mathcal{L}(V)$  be the set of all linearly independent subsets of  $V$  partially ordered by set containment.

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a totally ordered subset of  $\mathcal{L}(V)$ . That is for any sets  $\mathcal{B}_i, \mathcal{B}_j \in X$ , either  $\mathcal{B}_i \subseteq \mathcal{B}_j$  or  $\mathcal{B}_j \subseteq \mathcal{B}_i$ .

Let  $\mathcal{B} = \bigcup_{i \in I} \mathcal{B}_i$ . For all  $i \in I$ ,  $\mathcal{B}_i \subseteq \mathcal{B}$ . So  $\mathcal{B}$  is an upper bound for  $X$ .

**Claim 2:**  $\mathcal{B}$  is in  $\mathcal{L}(V)$ .

# Every vector space has a basis

Let  $\mathcal{L}(V)$  be the set of all linearly independent subsets of  $V$  partially ordered by set containment.

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a totally ordered subset of  $\mathcal{L}(V)$ . That is for any sets  $\mathcal{B}_i, \mathcal{B}_j \in X$ , either  $\mathcal{B}_i \subseteq \mathcal{B}_j$  or  $\mathcal{B}_j \subseteq \mathcal{B}_i$ .

Let  $\mathcal{B} = \bigcup_{i \in I} \mathcal{B}_i$ . For all  $i \in I$ ,  $\mathcal{B}_i \subseteq \mathcal{B}$ . So  $\mathcal{B}$  is an upper bound for  $X$ .

**Claim 2:**  $\mathcal{B}$  is in  $\mathcal{L}(V)$ .

**Zorn's Lemma:** Since every totally ordered subset of  $\mathcal{L}(V)$  has an upper bound in  $\mathcal{L}(V)$ ,  $\mathcal{L}(V)$  has a maximal element.

# Every vector space has a basis

Let  $\mathcal{L}(V)$  be the set of all linearly independent subsets of  $V$  partially ordered by set containment.

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a totally ordered subset of  $\mathcal{L}(V)$ . That is for any sets  $\mathcal{B}_i, \mathcal{B}_j \in X$ , either  $\mathcal{B}_i \subseteq \mathcal{B}_j$  or  $\mathcal{B}_j \subseteq \mathcal{B}_i$ .

Let  $\mathcal{B} = \bigcup_{i \in I} \mathcal{B}_i$ . For all  $i \in I$ ,  $\mathcal{B}_i \subseteq \mathcal{B}$ . So  $\mathcal{B}$  is an upper bound for  $X$ .

**Claim 2:**  $\mathcal{B}$  is in  $\mathcal{L}(V)$ .

**Zorn's Lemma:** Since every totally ordered subset of  $\mathcal{L}(V)$  has an upper bound in  $\mathcal{L}(V)$ ,  $\mathcal{L}(V)$  has a maximal element.

**Claim 1:** A maximal element of  $\mathcal{L}(V)$  is a basis for  $V$ .

# Every vector space has a basis

Let  $\mathcal{L}(V)$  be the set of all linearly independent subsets of  $V$  partially ordered by set containment.

Let  $X = \{\mathcal{B}_i \mid i \in I\}$  be a totally ordered subset of  $\mathcal{L}(V)$ . That is for any sets  $\mathcal{B}_i, \mathcal{B}_j \in X$ , either  $\mathcal{B}_i \subseteq \mathcal{B}_j$  or  $\mathcal{B}_j \subseteq \mathcal{B}_i$ .

Let  $\mathcal{B} = \bigcup_{i \in I} \mathcal{B}_i$ . For all  $i \in I$ ,  $\mathcal{B}_i \subseteq \mathcal{B}$ . So  $\mathcal{B}$  is an upper bound for  $X$ .

**Claim 2:**  $\mathcal{B}$  is in  $\mathcal{L}(V)$ .

**Zorn's Lemma:** Since every totally ordered subset of  $\mathcal{L}(V)$  has an upper bound in  $\mathcal{L}(V)$ ,  $\mathcal{L}(V)$  has a maximal element.

**Claim 1:** A maximal element of  $\mathcal{L}(V)$  is a basis for  $V$ .


Therefore,  $V$  has a basis.

# Connection: Graph Theory

A **graph** is a set of vertices and a set of edges joining some pairs of the vertices.



# Connection: Graph Theory

A **graph** is a set of vertices and a set of edges joining some pairs of the vertices.






# Connection: Graph Theory

A **graph** is a set of vertices and a set of edges joining some pairs of the vertices.






# Connection: Graph Theory

A **graph** is a set of vertices and a set of edges joining some pairs of the vertices.



# Connection: Graph Theory




A **graph** is a set of vertices and a set of edges joining some pairs of the vertices.



A graph is **connected** if there is a path from any vertex to any other vertex.

# Connection: Graph Theory

A **graph** is a set of vertices and a set of edges joining some pairs of the vertices.






A graph is **connected** if there is a path from any vertex to any other vertex.

A graph is a **tree** if there is a unique path from any vertex to any other vertex.

# Connection: Graph Theory

A **graph** is a set of vertices and a set of edges joining some pairs of the vertices.



A graph is **connected** if there is a path from any vertex to any other vertex.

A graph is a **tree** if there is a unique path from any vertex to any other vertex.

**Note:** A tree is a connected graph which does not contain any cycles.

# Connection: Graph Theory



A contradiction

# Connection: Graph Theory



A contradiction




Fife Council are good graph theorists.

# Connection: Graph Theory

A **spanning tree** of a graph is a subgraph which is a tree and which includes every vertex.


# Connection: Graph Theory

A **spanning tree** of a graph is a subgraph which is a tree and which includes every vertex.



# Connection: Graph Theory

A **spanning tree** of a graph is a subgraph which is a tree and which includes every vertex.




## Theorem

*Every connected graph has a spanning tree.*

# Connection: Graph Theory

A **spanning tree** of a graph is a subgraph which is a tree and which includes every vertex.



## Theorem


*Every connected graph has a spanning tree.*

## Sketch of Proof

Show that every maximal subtree is a spanning tree.

# Connection: Graph Theory

A **spanning tree** of a graph is a subgraph which is a tree and which includes every vertex.



## Theorem

*Every connected graph has a spanning tree.*


## Sketch of Proof

Show that every maximal subtree is a spanning tree.

Show that the union of any nest of trees is a tree.

# Connection: Graph Theory

A **spanning tree** of a graph is a subgraph which is a tree and which includes every vertex.



## Theorem

*Every connected graph has a spanning tree.*

## Sketch of Proof

Show that every maximal subtree is a spanning tree.

Show that the union of any nest of trees is a tree.

Conclude, by Zorn's Lemma, that every graph has a spanning tree. ■

# Closing Remarks

# Closing Remarks

Any questions?